

P416601WO PictaGip
Doppel

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 March 2002 (14.03.2002)

PCT

(10) International Publication Number
WO 02/20731 A2

(51) International Patent Classification?: C12N 9/00

(21) International Application Number: PCT/EP01/10125

(22) International Filing Date:
3 September 2001 (03.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/231,297 8 September 2000 (08.09.2000) US

(71) Applicant (for all designated States except US): BAYER
AKTIENGESELLSCHAFT [DE/DE]; 51368 Lev-
erkusen (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): XIAO, Yonghong
[US/US]; 78 Dana Street #1, Cambridge, MA 02138 (US).

(74) Common Representative: BAYER AKTIENGE-
SELLSCHAFT; 51368 Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: REGULATION OF HUMAN PROTEIN DISULFIDE ISOMERASE-LIKE ENZYME

(57) Abstract. Reagents which regulate human protein disulfide isomerase-like enzyme and reagents which bind to human protein disulfide isomerase-like enzyme gene products can play a role in preventing, ameliorating, or correcting dysfunctions or disorders including, but not limited to, cancer, diabetes, and obesity.

BEST AVAILABLE COPY

REGULATION OF HUMAN PROTEIN DISULFIDE ISOMERASE-LIKE ENZYME

5 TECHNICAL FIELD OF THE INVENTION

The invention relates to the area of regulation of human protein disulfide isomerase-like enzyme to provide therapeutic effects.

10 BACKGROUND OF THE INVENTION

Cells contain a number of proteins known as molecular chaperones or foldases. U.S. Patent 6,001,632. These molecules catalyze the folding of newly synthesized proteins, prevent aggregation and improper glycosylation, and remove denatured
15 proteins. Although they do not become part of the final structure, they are important in the assembly of proteins or their subunits into larger, more complex structures. In the absence of chaperones and foldases, misfolded proteins are quickly degraded by intracellular proteases.

20 Foldases, such as protein disulfide isomerase (PDI), are specialized enzymes which carry out rate-limiting covalent steps in protein folding. These enzymes are most abundant in cells actively synthesizing secreted proteins which are major components of the ER lumen. Tasanen *et al.*, *J. Biol. Chem.* 267, 11513-19, 1992) and may constitute 1-2% of eukaryotic cellular proteins. Although incubation of reduced
25 unfolded proteins in buffers with defined ratios of oxidized and reduced thiols can lead to native conformation, the rate of folding is slow and the attainment of native conformation decreases proportionately to the size and number of cysteines in the protein. In contrast, PDI in the eukaryotic ER is much more efficient in carrying out the enzymatic pairing and oxidation of cysteines.

30

Human PDI is a resident endoplasmic reticulum protein which mediates the formation of disulfide bonds in recently synthesized proteins. Chen *et al.*, *Br. J. Haematol.* 90, 425-31, 1995. PDI has recently been localized additional sites. Catalytically active PDI also has been localized to the plasma membrane of mammalian cells and platelets. PDI is also released from activated and inactivated platelets. *Id.* PDI catalyzes the reversible formation and isomerization of disulfide bonds, thus promoting correct protein folding. *Id.*; Essex *et al.*, *Blood* 86, 2168-73, 1995.

The activity of PDI is extremely concentration-dependent under certain conditions. PDI can promote the misfolding and aggregation of its substrates. When PDI is present at high concentrations relative to substrate, it acts a chaperone, increasing the yield of properly folded protein, and inhibiting substrate aggregation. In contrast when PDI is present at substoichiometric concentrations relative to substrate, it facilitates aggregation and inhibits proper protein folding. Sideraki & Gilbert, *Biochem.* 39, 1180-88, 2000.

PDI plays an important role in platelet responses. PDI has been shown to mediate platelet aggregation and secretion. PDI is further postulated to promote wound healing by facilitating the proper folding of proteins important for tissue remodeling. Essex & Li, *Br. J. Haematol.* 104, 448-54, 1999.

PDI also has been implicated tumor cell proliferation. Tumor cells responsive to the potent antiproliferative Tissue Growth Factor $\beta 1$ (TGF $\beta 1$) also down regulate PDI expression when treated with TGF $\beta 1$. TGF $\beta 1$ -resistant tumor cells, however do not alter their pattern of PDI expression after treatment with TGF $\beta 1$. PDI is postulated to interfere with TGF $\beta 1$ activity by either altering the folding of the TGF $\beta 1$ receptor or the TGF $\beta 1$ molecule itself. Sipes *et al.*, *Cell Growth & Diff.* 1, 241-46, 1990.

Because of the importance of PDI in a variety of cellular processes, there is a need in the art to identify additional related molecules which can be regulated to provide therapeutic effects.

5 **SUMMARY OF THE INVENTION**

It is an object of the invention to provide reagents and methods of regulating a human protein disulfide isomerase-like enzyme. This and other objects of the invention are provided by one or more of the embodiments described below.

10

One embodiment of the invention is a protein disulfide isomerase-like enzyme polypeptide comprising an amino acid sequence selected from the group consisting of:

15 amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:2,

amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:29,

20

the amino acid sequence shown in SEQ ID NO:2; and

the amino acid sequence shown in SEQ ID NO:29.

25 Yet another embodiment of the invention is a method of screening for agents which decrease extracellular matrix degradation. A test compound is contacted with a protein disulfide isomerase-like enzyme polypeptide comprising an amino acid sequence selected from the group consisting of:

30 amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:2,

amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:29,

5 the amino acid sequence shown in SEQ ID NO:2; and

the amino acid sequence shown in SEQ ID NO:29.

10 Binding between the test compound and the protein disulfide isomerase-like enzyme polypeptide is detected. A test compound which binds to the protein disulfide isomerase-like enzyme polypeptide is thereby identified as a potential agent for decreasing extracellular matrix degradation. The agent can work by decreasing the activity of the protein disulfide isomerase-like enzyme.

15 Another embodiment of the invention is a method of screening for agents which decrease extracellular matrix degradation. A test compound is contacted with a polynucleotide encoding a protein disulfide isomerase-like enzyme polypeptide, wherein the polynucleotide comprises a nucleotide sequence selected from the group consisting of:

20

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO:1,

25 nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO:28,

the nucleotide sequence shown in SEQ ID NO:1; and

the nucleotide sequence shown in SEQ ID NO:28.

30

Binding of the test compound to the polynucleotide is detected. A test compound which binds to the polynucleotide is identified as a potential agent for decreasing extracellular matrix degradation. The agent can work by decreasing the amount of the protein disulfide isomerase-like enzyme through interacting with the protein disulfide isomerase-like enzyme mRNA.

Another embodiment of the invention is a method of screening for agents which regulate extracellular matrix degradation. A test compound is contacted with a protein disulfide isomerase-like enzyme polypeptide comprising an amino acid sequence selected from the group consisting of:

amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:2,

amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:29,

the amino acid sequence shown in SEQ ID NO:2; and

the amino acid sequence shown in SEQ ID NO:29.

A protein disulfide isomerase-like enzyme activity of the polypeptide is detected. A test compound which increases protein disulfide isomerase-like enzyme activity of the polypeptide relative to protein disulfide isomerase-like enzyme activity in the absence of the test compound is thereby identified as a potential agent for increasing extracellular matrix degradation. A test compound which decreases protein disulfide isomerase-like enzyme activity of the polypeptide relative to protein disulfide isomerase-like enzyme activity in the absence of the test compound is thereby identified as a potential agent for decreasing extracellular matrix degradation.

30

- 6 -

Even another embodiment of the invention is a method of screening for agents which decrease extracellular matrix degradation. A test compound is contacted with a protein disulfide isomerase-like enzyme product of a polynucleotide which comprises a nucleotide sequence selected from the group consisting of:

5

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO:1,

10

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO:28,

the nucleotide sequence shown in SEQ ID NO:1; and

15

the nucleotide sequence shown in SEQ ID NO:28.

Binding of the test compound to the protein disulfide isomerase-like enzyme product is detected. A test compound which binds to the protein disulfide isomerase-like enzyme product is thereby identified as a potential agent for decreasing extracellular matrix degradation.

20

Still another embodiment of the invention is a method of reducing extracellular matrix degradation. A cell is contacted with a reagent which specifically binds to a polynucleotide encoding a protein disulfide isomerase-like enzyme polypeptide or the product encoded by the polynucleotide, wherein the polynucleotide comprises a nucleotide sequence selected from the group consisting of:

25

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO:1,

30

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO:28,

the nucleotide sequence shown in SEQ ID NO:1; and

the nucleotide sequence shown in SEQ ID NO:28.

5

Protein disulfide isomerase-like enzyme activity in the cell is thereby decreased.

The invention thus provides a human protein disulfide isomerase-like enzyme which can be used to identify test compounds which may act, for example, as agonists or antagonists at the enzyme's active site. Human protein disulfide isomerase-like enzyme and fragments thereof also are useful in raising specific antibodies which can block the enzyme and effectively reduce its activity.

10

BRIEF DESCRIPTION OF THE DRAWINGS

15

Fig. 1 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:1).

20

Fig. 2 shows the amino acid sequence deduced from the DNA-sequence of Fig.1 (SEQ ID NO:2).

Fig. 3 shows the amino acid sequence of the protein identified by SwissProt Accession No. Q10057 (SEQ ID NO:3).

25

Fig. 4 shows the amino acid sequence of pfam/hmm/thioredoxin (SEQ ID NO:4).

Fig. 5 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:5).

30

Fig. 6 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:6).

Fig. 7 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:7).

5 Fig. 8 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:8).

Fig. 9 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:9).

10 Fig. 10 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:10).

15 Fig. 11 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:11).

Fig. 12 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:12).

20 Fig. 13 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:13).

Fig. 14 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:14).

25 Fig. 15 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:15).

30 Fig. 16 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:16).

- Fig. 17 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:17).
- 5 Fig. 18 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:18).
- Fig. 19 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:19).
- 10 Fig. 20 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:20).
- Fig. 21 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:21).
- 15 Fig. 22 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:22).
- Fig. 23 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:23).
- 20 Fig. 24 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:24).
- Fig. 25 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:25).
- 25 Fig. 26 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:26).
- 30

Fig. 27 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:27).

5 Fig. 28 shows the BLASTP alignment of 59_PROTEIN (SEQ ID NO:2) against swiss|Q10057|PDI1_SCHPO (SEQ ID NO:3).

Fig. 29 shows the BLOCKS search results.

10 Fig. 30 shows the HMMPFAM - alignment of 59_PROTEIN (SEQ ID NO:2) against pfam|hmm|thiorel (SEQ ID NO:4)

Fig. 31 shows the DNA-sequence encoding a protein disulfide isomerase-like enzyme polypeptide (SEQ ID NO:28).

15 Fig. 32 shows the amino acid sequence deduced from the DNA-sequence of Fig. 31 (SEQ ID NO:29).

Fig. 33 shows the BLASTP alignment of 59_short_TR1 against swiss/P30101/ER60_HUMAN.

20

Fig. 34 shows the HMMPFAM alignment of 59_short_TR1 against pfam/hmm/thiorel.

25 Fig. 35 shows the expression profile of protein disulfide isomerase-like protein mRNA in whole body tissues.

Fig. 36 shows the expression profile of protein disulfide isomerase-like protein mRNA in blood and lung tissues.

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to an isolated polynucleotide encoding a protein disulfide isomerase-like enzyme polypeptide and being selected from the group consisting of:

5

- a) a polynucleotide encoding a protein disulfide isomerase-like enzyme polypeptide comprising an amino acid sequence selected from the group consisting of:

10

amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:2,
amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:29,

15

the amino acid sequence shown in SEQ ID NO:2; and
the amino acid sequence shown in SEQ ID NO:29.

20

- b) a polynucleotide comprising the sequence of SEQ ID NO:1 or SEQ ID NO:28;

- c) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) and (b);

25

- d) a polynucleotide the sequence of which deviates from the polynucleotide sequences specified in (a) to (c) due to the degeneration of the genetic code; and

30

- e) a polynucleotide which represents a fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (d).

- 12 -

Furthermore, it has been discovered by the present applicant that a novel protein disulfide isomerase-like enzyme, particularly a human protein disulfide isomerase-like enzyme, is a discovery of the present invention. Human protein disulfide isomerase-like enzyme comprises the amino acid sequences shown in SEQ ID NO:2 and SEQ ID NO:29, as encoded by the coding sequence shown in SEQ ID NO:1 and SEQ ID NO:28, respectively.

Human protein disulfide isomerase-like enzyme is 32% identical over a 131 amino acid overlap to the protein identified by SwissProt Accession No. Q10057 (SEQ ID NO:3) and annotated as "PUTATIVE PROTEIN DISULFIDE ISOMERASE C1F5.02 PRECURSOR (EC 5.3.4.1)." (FIG. 28).

The human protein disulfide isomerase-like enzyme of the invention is expected to be useful for the same purposes as previously identified protein disulfide isomerase enzymes. For example, protein disulfide isomerase is found on the surface of platelets and is released upon platelet activation and mediates platelet aggregation and secretion. Because platelets are present and become activated when and where vascular injury occurs, platelet protein disulfide isomerase and, therefore, human protein disulfide isomerase-like enzyme, may play a role in the various hemostatic and tissue remodeling processes in which platelets (hence blood coagulation) are involved. Protein disulfide isomerase also has a regulatory effect on the surface protein thiol status of B cells. The increased expression of protein disulfide isomerase may play a crucial role in SH-mediated protection and drug resistance in malignant B lymphocytes. Protein disulfide isomerase mRNA levels correlate inversely with sensitivity to the growth-inhibitory effects of TGF β 1, and protein disulfide isomerase down-regulation occurs in some TGF β -sensitive cells, but not in a human pancreatic carcinoma cell line which is insensitive to the growth-inhibitory effects of TGF β 1.

Because human protein disulfide isomerase-like enzyme is expressed mainly in tumor and other highly proliferative tissues (*i.e.*, tissues with a rich blood supply), it

- 13 -

is a potential target for diseases therapeutic sensitivity/resistance involving blood clotting, vascular wall repair and formation, as well as tissue remodeling. Human protein disulfide isomerase-like enzyme therefore can be used in therapeutic methods to treat disorders such as hematological disorders, cardiovascular disorders, and cancer. Human protein disulfide isomerase-like enzyme also can be used to screen for human protein disulfide isomerase-like enzyme agonists and antagonists.

Polypeptides

Protein disulfide isomerase-like polypeptides according to the invention comprise at least 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, or 390 contiguous amino acids selected from the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29 or a biologically active variant thereof, as defined below. A protein disulfide isomerase-like polypeptide of the invention therefore can be a portion of a protein disulfide isomerase-like enzyme, a full-length protein disulfide isomerase-like enzyme, or a fusion protein comprising all or a portion of a protein disulfide isomerase-like enzyme.

Biologically Active Variants

Protein disulfide isomerase-like polypeptide variants which are biologically active, *i.e.*, retain a protein disulfide isomerase-like activity, also are protein disulfide isomerase-like polypeptides. Preferably, naturally or non-naturally occurring protein disulfide isomerase-like polypeptide variants have amino acid sequences which are at least about 50, 55, 60, 65, or 70, preferably about 75, 80, 85, 90, 96, 96, or 98% identical to the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29 or a fragment thereof. Percent identity between a putative protein disulfide isomerase-like polypeptide variant and an amino acid sequence of SEQ ID NO:2 or SEQ ID NO:29 is determined using the Blast2 alignment program (Blosum62, Expect 10, standard genetic codes).

Variations in percent identity can be due, for example, to amino acid substitutions, insertions, or deletions. Amino acid substitutions are defined as one for one amino acid replacements. They are conservative in nature when the substituted amino acid has similar structural and/or chemical properties. Examples of conservative replacements are substitution of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

Amino acid insertions or deletions are changes to or within an amino acid sequence. They typically fall in the range of about 1 to 5 amino acids. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological or immunological activity of a protein disulfide isomerase-like polypeptide can be found using computer programs well known in the art, such as DNASTAR software. Whether an amino acid change results in a biologically active protein disulfide isomerase-like polypeptide can readily be determined by assaying for protein disulfide isomerase-like activity, as described for example, in Sideraki & Gilbert, 2000, Essex *et al.*, 1995, or Chen *et al.*, 1992.

Fusion Proteins

Fusion proteins are useful for generating antibodies against protein disulfide isomerase-like polypeptide amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of a protein disulfide isomerase-like polypeptide. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

A protein disulfide isomerase-like polypeptide fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment comprises at least 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, or 390 contiguous amino acids of SEQ ID NO:2 or SEQ ID NO:29 or

of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length protein disulfide isomerase-like enzyme protein.

- 5 The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include β -galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horse-radish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Addi-
- 10 tionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A
- 15 fusion protein also can be engineered to contain a cleavage site located between the protein disulfide isomerase-like polypeptide-encoding sequence and the heterologous protein sequence, so that the protein disulfide isomerase-like polypeptide can be cleaved and purified away from the heterologous moiety.
- 20 A fusion protein can be synthesized chemically, as is known in the art. Preferably, a fusion protein is produced by covalently linking two polypeptide segments or by standard procedures in the art of molecular biology. Recombinant DNA methods can be used to prepare fusion proteins, for example, by making a DNA construct which comprises coding sequences selected from SEQ ID NO:1 or SEQ ID NO:28 in
- 25 proper reading frame with nucleotides encoding the second polypeptide segment and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies such as Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), CLONTECH (Mountain View, CA), Santa Cruz Biotechnology (Santa Cruz, CA), MBL International
- 30 Corporation (MIC; Watertown, MA), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

Identification of Species Homologs

Species homologs of human protein disulfide isomerase-like polypeptide can be obtained using protein disulfide isomerase-like polypeptide polynucleotides (described below) to make suitable probes or primers for screening cDNA expression libraries from other species, such as mice, monkeys, or yeast, identifying cDNAs which encode homologs of protein disulfide isomerase-like polypeptide, and expressing the cDNAs as is known in the art.

Polynucleotides

A protein disulfide isomerase-like polynucleotide can be single- or double-stranded and comprises a coding sequence or the complement of a coding sequence for a protein disulfide isomerase-like polypeptide. A coding sequence for human protein disulfide isomerase-like enzyme is shown in SEQ ID NO:1 and SEQ ID NO:28. This coding sequence is found within the larger genomic sequence shown in SEQ ID NO:5.

Degenerate nucleotide sequences encoding human protein disulfide isomerase-like polypeptides, as well as homologous nucleotide sequences which are at least about 50, 55, 60, 65, 70, preferably about 75, 90, 96, or 98% identical to the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:28 also are protein disulfide isomerase-like polynucleotides. Percent sequence identity between the sequences of two polynucleotides is determined using computer programs such as ALIGN which employ the FASTA algorithm, using an affine gap search with a gap open penalty of -12 and a gap extension penalty of -2. Complementary DNA (cDNA) molecules, species homologs, and variants of protein disulfide isomerase-like polynucleotides which encode biologically active protein disulfide isomerase-like polypeptides also are protein disulfide isomerase-like polynucleotides.

Identification of Polynucleotide Variants and Homologs

5 Variants and homologs of the protein disulfide isomerase-like polynucleotides described above also are protein disulfide isomerase-like polynucleotides. Typically, homologous protein disulfide isomerase-like polynucleotide sequences can be identified by hybridization of candidate polynucleotides to known protein disulfide isomerase-like polynucleotides under stringent conditions, as is known in the art. For example, using the following wash conditions: 2X SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7.0), 0.1% SDS, room temperature twice, 30 minutes each; then 10 2X SSC, 0.1% SDS, 50 EC once, 30 minutes; then 2X SSC, room temperature twice, 10 minutes each homologous sequences can be identified which contain at most about 25-30% basepair mismatches. More preferably, homologous nucleic acid strands contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches.

15 Species homologs of the protein disulfide isomerase-like polynucleotides disclosed herein also can be identified by making suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, or yeast. Human variants of protein disulfide isomerase-like polynucleotides can be identified, 20 for example, by screening human cDNA expression libraries. It is well known that the T_m of a double-stranded DNA decreases by 1-1.5°C with every 1% decrease in homology (Bonner *et al.*, *J. Mol. Biol.* 81, 123 (1973). Variants of human protein disulfide isomerase-like polynucleotides or protein disulfide isomerase-like polynucleotides of other species can therefore be identified by hybridizing a putative 25 homologous protein disulfide isomerase-like polynucleotide with a polynucleotide having a nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:28 or the complement thereof to form a test hybrid. The melting temperature of the test hybrid is compared with the melting temperature of a hybrid comprising polynucleotides having perfectly complementary nucleotide sequences, and the number or percent of 30 basepair mismatches within the test hybrid is calculated.

- 18 -

Nucleotide sequences which hybridize to protein disulfide isomerase-like polynucleotides or their complements following stringent hybridization and/or wash conditions also are protein disulfide isomerase-like polynucleotides. Stringent wash conditions are well known and understood in the art and are disclosed, for example, in Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2d ed., 1989, at pages 9.50-9.51.

Typically, for stringent hybridization conditions a combination of temperature and salt concentration should be chosen that is approximately 12-20°C below the calculated T_m of the hybrid under study. The T_m of a hybrid between a protein disulfide isomerase-like polynucleotide having a nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:28 or the complement thereof and a polynucleotide sequence which is at least about 50, 55, 60, 65, 70, preferably about 75, 90, 96, or 98% identical to one of those nucleotide sequences can be calculated, for example, using the equation of Bolton and McCarthy, *Proc. Natl. Acad. Sci. U.S.A.* 48, 1390 (1962):

$$T_m = 81.5^{\circ}\text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\%G + C) - 0.63(\%\text{formamide}) - 600/l,$$

where l = the length of the hybrid in basepairs.

Stringent wash conditions include, for example, 4X SSC at 65°C, or 50% formamide, 4X SSC at 42°C, or 0.5X SSC, 0.1% SDS at 65°C. Highly stringent wash conditions include, for example, 0.2X SSC at 65°C.

Preparation of Polynucleotides

A naturally occurring protein disulfide isomerase-like polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such

- 19 -

technique for obtaining a polynucleotide can be used to obtain isolated protein disulfide isomerase-like polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprises protein disulfide isomerase-like enzyme nucleotide sequences. Isolated polynucleotides are
5 in preparations which are free or at least 70, 80, or 90% free of other molecules.

Protein disulfide isomerase-like enzyme cDNA molecules can be made with standard molecular biology techniques, using protein disulfide isomerase-like enzyme mRNA as a template. Protein disulfide isomerase-like enzyme cDNA molecules can
10 thereafter be replicated using molecular biology techniques known in the art and disclosed in manuals such as Sambrook *et al.* (1989). An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

15 Alternatively, synthetic chemistry techniques can be used to synthesize protein disulfide isomerase-like polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode a protein disulfide isomerase-like polypeptide having, for example, an amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29 or a biologically active variant thereof.

20

Extending Polynucleotides

The partial nucleotide sequence disclosed herein can be used to identify the corresponding full length gene from which they were derived. The partial sequence
25 can be nick-translated or end-labeled with ³²P using polynucleotide kinase using labeling methods known to those with skill in the art (BASIC METHODS IN MOLECULAR BIOLOGY, Davis *et al.*, eds., Elsevier Press, N.Y., 1986). A lambda library prepared from human tissue can be directly screened with the labeled sequences of interest or the library can be converted en masse to pBluescript
30 (Stratagene Cloning Systems, La Jolla, Calif. 92037) to facilitate bacterial colony

- 20 -

screening (see Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press (1989, pg. 1.20).

Both methods are well known in the art. Briefly, filters with bacterial colonies
5 containing the library in pBluescript or bacterial lawns containing lambda plaques are denatured, and the DNA is fixed to the filters. The filters are hybridized with the labeled probe using hybridization conditions described by Davis *et al.*, 1986. The partial sequences, cloned into lambda or pBluescript, can be used as positive controls to assess background binding and to adjust the hybridization and washing
10 stringencies necessary for accurate clone identification. The resulting autoradiograms are compared to duplicate plates of colonies or plaques; each exposed spot corresponds to a positive colony or plaque. The colonies or plaques are selected, expanded and the DNA is isolated from the colonies for further analysis and sequencing.

15 Positive cDNA clones are analyzed to determine the amount of additional sequence they contain using PCR with one primer from the partial sequence and the other primer from the vector. Clones with a larger vector-insert PCR product than the original partial sequence are analyzed by restriction digestion and DNA sequencing
20 to determine whether they contain an insert of the same size or similar as the mRNA size determined from Northern blot Analysis.

Once one or more overlapping cDNA clones are identified, the complete sequence of the clones can be determined, for example after exonuclease III digestion
25 (McCombie *et al.*, *Methods* 3, 33-40, 1991). A series of deletion clones are generated, each of which is sequenced. The resulting overlapping sequences are assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a highly accurate final sequence.

30

- 21 -

Various PCR-based methods can be used to extend the nucleic acid sequences disclosed herein to detect upstream sequences such as promoters and regulatory elements. For example, restriction site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, *PCR Methods Applic.* 2, 318322, 1993). Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region (Triglia *et al.*, *Nucleic Acids Res.* 16, 8186, 1988). Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 2230 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA (Lagerstrom *et al.*, *PCR Methods Applic.* 1, 111119, 1991). In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

Another method which can be used to retrieve unknown sequences is that of Parker *et al.*, *Nucleic Acids Res.* 19, 30553060, 1991). Additionally, PCR, nested primers, and PROMOTERFINDER libraries (CLONTECH, Palo Alto, Calif.) can be used to

- 22 -

walk genomic DNA (CLONTECH, Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

5 When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' nontranscribed regulatory regions.

10

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser
15 activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate software (e.g. GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable
20 for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

Obtaining Polypeptides

25 Protein disulfide isomerase-like polypeptides can be obtained, for example, by purification from human cells, by expression of protein disulfide isomerase-like polynucleotides, or by direct chemical synthesis.

Protein Purification

Protein disulfide isomerase-like polypeptides can be purified from any cell which expresses the enzyme, including host cells which have been transfected with protein

5 disulfide isomerase-like enzyme expression constructs. Testis, colon tumor, lung tumor, neuroendocrine lung carcinoid, glioblastoma, immortalized keratinocytes, normal lung, fetal liver, spleen, pancreatic adenocarcinoma, normal prostate, germinal center B cells, germ cell tumors, papillary serous carcinoma, breast tumor, kidney tumors, infant brain cDNA from a muscular atrophy patient, multiple

10 sclerosis lesions, fetal brain, uterus, Burkitt's lymphoma, and choriocarcinoma provide useful sources of protein disulfide isomerase-like enzyme. A purified protein disulfide isomerase-like polypeptide is separated from other compounds which normally associate with the protein disulfide isomerase-like polypeptide in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the

15 art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis. A preparation of purified protein disulfide isomerase-like polypeptides is at least 80% pure; preferably, the preparations are 90%, 95%, or 99% pure. Purity of the preparations can be assessed

20 by any means known in the art, such as SDS-polyacrylamide gel electrophoresis.

Expression of Polynucleotides

To express a protein disulfide isomerase-like polynucleotide, the polynucleotide can

25 be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding protein disulfide isomerase-like polypeptides and appropriate transcriptional and translational control elements. These methods

30 include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. Such techniques are described, for example, in Sambrook *et*

- 24 -

al. (1989) and in Ausubel *et al.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, N.Y., 1989.

A variety of expression vector/host systems can be utilized to contain and express sequences encoding a protein disulfide isomerase-like polypeptide. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (*e.g.*, baculovirus), plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids), or animal cell systems.

The control elements or regulatory sequences are those nontranslated regions of the vector enhancers, promoters, 5' and 3' untranslated regions which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (*e.g.*, heat shock, RUBISCO, and storage protein genes) or from plant viruses (*e.g.*, viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding a protein disulfide isomerase-like polypeptide, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the protein disulfide isomerase-like polypeptide. For example, when a large quantity of a protein disulfide isomerase-like polypeptide is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding the protein disulfide isomerase-like polypeptide can be ligated into the vector in frame with sequences for the amino terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors (Van Heeke & Schuster, *J. Biol. Chem.* 264, 55035509, 1989) or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH can be used. For reviews, see Ausubel *et al.* (1989) and Grant *et al.*, *Methods Enzymol.* 153, 516544, 1987.

Plant and Insect Expression Systems

If plant expression vectors are used, the expression of sequences encoding protein disulfide isomerase-like polypeptides can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be

- 26 -

used alone or in combination with the omega leader sequence from TMV (Takamatsu, *EMBO J.* 6, 307311, 1987). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used (Coruzzi *et al.*, *EMBO J.* 3, 16711680, 1984; Broglie *et al.*, *Science* 224, 838843, 1984; Winter *et al.*, *Results Probl. Cell Differ.* 17, 85105, 1991). These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (*e.g.*, Hobbs or Murray, in MCGRAW HILL YEARBOOK OF SCIENCE AND TECHNOLOGY, McGraw Hill, New York, N.Y., pp. 191196, 1992).

An insect system also can be used to express a protein disulfide isomerase-like polypeptide. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding protein disulfide isomerase-like polypeptides can be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of protein disulfide isomerase-like polypeptides will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which protein disulfide isomerase-like polypeptides can be expressed (Engelhard *et al.*, *Proc. Nat. Acad. Sci.* 91, 32243227, 1994).

Mammalian Expression Systems

A number of viral-based expression systems can be used to express protein disulfide isomerase-like polypeptides in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding protein disulfide isomerase-like polypeptides can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing a protein disulfide isomerase-like polypeptide in

- 27 -

infected host cells (Logan & Shenk, *Proc. Natl. Acad. Sci.* 81, 36553659, 1984). If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

- 5 Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (*e.g.*, liposomes, polycationic amino polymers, or vesicles).
- 10 Specific initiation signals also can be used to achieve more efficient translation of sequences encoding protein disulfide isomerase-like polypeptides. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding a protein disulfide isomerase-like polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional
- 15 transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of
- 20 various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used (see Scharf *et al.*, *Results Probl. Cell Differ.* 20, 125162, 1994).

Host Cells

- 25 A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein disulfide isomerase-like polypeptide in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation,
- 30 lipidation, and acylation. Posttranslational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or

- 28 -

function. Different host cells which have specific cellular machinery and characteristic mechanisms for Post-translational activities (*e.g.*, CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen
5 to ensure the correct modification and processing of the foreign protein.

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express protein disulfide isomerase-like polypeptides can be transformed using expression vectors which can contain
10 viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 12 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which
15 successfully express the introduced protein disulfide isomerase-like enzyme sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. See, for example, ANIMAL CELL CULTURE, R.I. Freshney, ed., 1986.

20 Any number of selection systems can be used to recover transformed cell lines.

These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler *et al.*, *Cell* 11, 22332, 1977) and adenine phosphoribosyltransferase (Lowy *et al.*, *Cell* 22, 81723, 1980) genes which can be employed in *tk*⁻ or *aprt*⁻ cells,
25 respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate (Wigler *et al.*, *Proc. Natl. Acad. Sci.* 77, 356770, 1980), *npt* confers resistance to the aminoglycosides, neomycin and G418 (Colbere-Garapin *et al.*, *J. Mol. Biol.* 150, 114, 1981), and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin
30 acetyltransferase, respectively (Murray, 1992, *supra*). Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of

tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci.* 85, 804751, 1988). Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes *et al.*, *Methods Mol. Biol.* 55, 121131, 1995).

Detecting Expression

Although the presence of marker gene expression suggests that the protein disulfide isomerase-like polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding a protein disulfide isomerase-like polypeptide is inserted within a marker gene sequence, transformed cells containing sequences which encode a protein disulfide isomerase-like polypeptide can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a protein disulfide isomerase-like polypeptide under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the protein disulfide isomerase-like polynucleotide.

Alternatively, host cells which contain a protein disulfide isomerase-like polynucleotide and which express a protein disulfide isomerase-like polypeptide can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding a protein disulfide isomerase-like polypeptide can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding a protein disulfide isomerase-like polypeptide. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from

- 30 -

sequences encoding a protein disulfide isomerase-like polypeptide to detect transformants which contain a protein disulfide isomerase-like polynucleotide.

5 A variety of protocols for detecting and measuring the expression of a protein disulfide isomerase-like polypeptide, using either polyclonal or monoclonal anti-
bodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence
10 activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on a protein disulfide isomerase-like polypeptide can be used, or a competitive binding assay can be employed. These and other assays are described in Hampton *et al.*, SEROLOGICAL METHODS: A LABORATORY MANUAL, APS Press, St. Paul, Minn., 1990) and Maddox *et al.*, *J. Exp. Med.* 158, 12111216, 1983).

15 A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding protein disulfide isomerase-like polypeptides include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled
20 nucleotide. Alternatively, sequences encoding a protein disulfide isomerase-like polypeptide can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes *in vitro* by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using
25 a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

30

Expression and Purification of Polypeptides

Host cells transformed with nucleotide sequences encoding a protein disulfide isomerase-like polypeptide can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode protein disulfide isomerase-like polypeptides can be designed to contain signal sequences which direct secretion of soluble protein disulfide isomerase-like polypeptides through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound protein disulfide isomerase-like polypeptide.

As discussed above, other constructions can be used to join a sequence encoding a protein disulfide isomerase-like polypeptide to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor Xa or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the protein disulfide isomerase-like polypeptide also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a protein disulfide isomerase-like polypeptide and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography, as described in Porath *et al.*, *Prot. Exp. Purif.* 3, 263281, 1992), while the enterokinase cleavage site provides a means for purifying the protein disulfide isomerase-like polypeptide from the fusion protein. Vectors which contain fusion proteins are disclosed in Kroll *et al.*, *DNA Cell Biol.* 12, 441453, 1993.

Chemical Synthesis

Sequences encoding a protein disulfide isomerase-like polypeptide can be synthesized, in whole or in part, using chemical methods well known in the art (see
5 Caruthers *et al.*, *Nucl. Acids Res. Symp. Ser.* 215223, 1980; Horn *et al.* *Nucl. Acids Res. Symp. Ser.* 225232, 1980). Alternatively, a protein disulfide isomerase-like polypeptide itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques
10 (Merrifield, *J. Am. Chem. Soc.* 85, 21492154, 1963; Roberge *et al.*, *Science* 269, 202204, 1995). Protein synthesis can be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of protein disulfide isomerase-like polypeptides can be separately synthesized and
15 combined using chemical methods to produce a full-length molecule.

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography (*e.g.*, Creighton, *PROTEINS: STRUCTURES AND MOLECULAR PRINCIPLES*, WH Freeman and Co., New York, N.Y., 1983). The
20 composition of a synthetic protein disulfide isomerase-like polypeptide can be confirmed by amino acid analysis or sequencing (*e.g.*, the Edman degradation procedure; *see* Creighton, *supra*). Additionally, any portion of the amino acid sequence of the protein disulfide isomerase-like polypeptide can be altered during direct synthesis and/or combined using chemical methods with sequences from other
25 proteins to produce a variant polypeptide or a fusion protein.

Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce
30 protein disulfide isomerase-like polypeptide-encoding nucleotide sequences possessing non-natural occurring codons. For example, codons preferred by a

particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

5

The nucleotide sequences disclosed herein can be engineered using methods generally known in the art to alter protein disulfide isomerase-like polypeptide-encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

15

Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of a protein disulfide isomerase-like polypeptide. "Antibody" as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')₂, and Fv, which are capable of binding an epitope of a protein disulfide isomerase-like polypeptide. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, *e.g.*, at least 15, 25, or 50 amino acids.

25

An antibody which specifically binds to an epitope of a protein disulfide isomerase-like polypeptide can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well

30

known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the immunogen.

5 Typically, an antibody which specifically binds to a protein disulfide isomerase-like polypeptide provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to protein disulfide isomerase-like polypeptides do not detect other proteins in immunochemical assays and can
10 immunoprecipitate a protein disulfide isomerase-like polypeptide from solution.

Protein disulfide isomerase-like polypeptides can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, a protein disulfide isomerase-like polypeptide can be
15 conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (*e.g.*, aluminum hydroxide), and surface active substances (*e.g.* lysolecithin, pluronic polyols, polyanions, peptides, oil
20 emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are especially useful.

Monoclonal antibodies which specifically bind to a protein disulfide isomerase-like
25 polypeptide can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B cell hybridoma technique, and the EBV hybridoma technique (Kohler *et al.*, *Nature* 256, 495497, 1985; Kozbor *et al.*, *J. Immunol. Methods* 81, 3142, 1985; Cote *et al.*, *Proc. Natl. Acad. Sci.* 80, 20262030, 1983; Cole *et al.*, *Mol. Cell Biol.* 62, 109120, 1984).
30

- 35 -

In addition, techniques developed for the production of "chimeric antibodies," the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used (Morrison *et al.*, *Proc. Natl. Acad. Sci.* 81, 68516855, 1984; Neuberger *et al.*, *Nature* 312, 604608, 1984; Takeda *et al.*, *Nature* 314, 452454, 1985). Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Alternatively, humanized antibodies can be produced using recombinant methods, as described in GB2188638B. Antibodies which specifically bind to a protein disulfide isomerase-like polypeptide can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. Patent 5,565,332.

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to protein disulfide isomerase-like polypeptides. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton, *Proc. Natl. Acad. Sci.* 88, 1112023, 1991).

Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template (Thirion *et al.*, 1996, *Eur. J. Cancer Prev.* 5, 507-11). Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in Coloma & Morrison, 1997, *Nat. Biotechnol.* 15, 159-63. Construction of bivalent, bispecific single-chain antibodies is taught in Mallender & Voss, 1994, *J. Biol. Chem.* 269, 199-206.

A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology (Verhaar *et al.*, 1995, *Int. J. Cancer* 61, 497-501; Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165, 81-91).

Antibodies which specifically bind to protein disulfide isomerase-like polypeptides also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi *et al.*, *Proc. Natl. Acad. Sci.* 86, 3833-3837, 1989; Winter *et al.*, *Nature* 349, 293-299, 1991).

Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which a protein disulfide isomerase-like polypeptide is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes

- 37 -

and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of protein disulfide isomerase-like enzyme gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamides, carboxymethyl esters, carbonates, and phosphate triesters. See Brown, *Meth. Mol. Biol.* 20, 18, 1994; Sonveaux, *Meth. Mol. Biol.* 26, 1-72, 1994; Uhlmann *et al.*, *Chem. Rev.* 90, 543583, 1990.

Modifications of protein disulfide isomerase-like enzyme gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the protein disulfide isomerase-like enzyme gene. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions 10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature (*e.g.*, Gee *et al.*, in Huber & Carr, *MOLECULAR AND IMMUNOLOGIC APPROACHES*, Futura Publishing Co., Mt. Kisco, N.Y., 1994). An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

30

- 38 -

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a protein disulfide isomerase-like polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a protein disulfide isomerase-like polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent protein disulfide isomerase-like enzyme nucleotides, can provide sufficient targeting specificity for protein disulfide isomerase-like enzyme mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular protein disulfide isomerase-like polynucleotide sequence.

Antisense oligonucleotides can be modified without affecting their ability to hybridize to a protein disulfide isomerase-like polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5' substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art. *See, e.g., Agrawal et al., Trends Biotechnol. 10, 152158, 1992; Uhlmann et al., Chem. Rev. 90, 543584, 1990; Uhlmann et al., Tetrahedron. Lett. 215, 35393542, 1987.*

Ribozymes

Ribozymes are RNA molecules with catalytic activity. *See, e.g., Cech, Science 236,*
15321539; 1987; Cech, *Ann. Rev. Biochem.* 59, 543568; 1990, Cech, *Curr. Opin.*
5 *Struct. Biol.* 2, 605609; 1992, Couture & Stinchcomb, *Trends Genet.* 12, 510515,
1996. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence,
as is known in the art (*e.g., Haseloff et al., U.S. Patent 5,641,673*). The mechanism
of ribozyme action involves sequence-specific hybridization of the ribozyme
molecule to complementary target RNA, followed by endonucleolytic cleavage.
10 Examples include engineered hammerhead motif ribozyme molecules that can
specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide
sequences.

The coding sequence of a protein disulfide isomerase-like polynucleotide can be used
15 to generate ribozymes which will specifically bind to mRNA transcribed from the
protein disulfide isomerase-like polynucleotide. Methods of designing and
constructing ribozymes which can cleave other RNA molecules in trans in a highly
sequence specific manner have been developed and described in the art (*see Haseloff*
et al. Nature 334, 585591, 1988). For example, the cleavage activity of ribozymes
20 can be targeted to specific RNAs by engineering a discrete "hybridization" region
into the ribozyme. The hybridization region contains a sequence complementary to
the target RNA and thus specifically hybridizes with the target (*see, for example,*
Gerlach et al., EP 321,201).

Specific ribozyme cleavage sites within a protein disulfide isomerase-like enzyme
25 RNA target can be identified by scanning the target molecule for ribozyme cleavage
sites which include the following sequences: GUA, GUU, and GUC. Once
identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding
to the region of the target RNA containing the cleavage site can be evaluated for
30 secondary structural features which may render the target inoperable. Suitability of
candidate protein disulfide isomerase-like enzyme RNA targets also can be evaluated

- 40 -

by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease protein disulfide isomerase-like enzyme expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells.

As taught in Haseloff *et al.*, U.S. Patent 5,641,673, ribozymes can be engineered so that ribozyme expression will occur in response to factors which induce expression of a target gene. Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

Differentially Expressed Genes

Described herein are methods for the identification of genes whose products interact with human protein disulfide isomerase-like enzyme. Such genes may represent genes which are differentially expressed in disorders including, but not limited to, cancer, diabetes, and obesity. Further, such genes may represent genes which are

- 41 -

differentially regulated in response to manipulations relevant to the progression or treatment of such diseases. Additionally, such genes may have a temporally modulated expression, increased or decreased at different stages of tissue or organism development. A differentially expressed gene may also have its expression modulated under control versus experimental conditions. In addition, the human protein disulfide isomerase-like gene or gene product may itself be tested for differential expression.

The degree to which expression differs in a normal versus a diseased state need only be large enough to be visualized via standard characterization techniques such as differential display techniques. Other such standard characterization techniques by which expression differences may be visualized include but are not limited to, quantitative RT (reverse transcriptase), PCR, and Northern analysis.

Identification of Differentially Expressed Genes

To identify differentially expressed genes total RNA or, preferably, mRNA is isolated from tissues of interest. For example, RNA samples are obtained from tissues of experimental subjects and from corresponding tissues of control subjects. Any RNA isolation technique which does not select against the isolation of mRNA may be utilized for the purification of such RNA samples. See, for example, Ausubel *et al.*, ed.,, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, Inc. New York, 1987-1993. Large numbers of tissue samples may readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski, U.S. Patent 4,843,155.

Transcripts within the collected RNA samples which represent RNA produced by differentially expressed genes are identified by methods well known to those of skill in the art. They include, for example, differential screening (Tedder *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 85, 208-12, 1988), subtractive hybridization (Hedrick *et al.*, *Nature* 308, 149-53; Lee *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 88, 2825, 1984), and,

preferably, differential display (Liang & Pardee, *Science* 257, 967-71, 1992; U.S. Patent 5,262,311).

5 The differential expression information may itself suggest relevant methods for the treatment of disorders involving the human protein disulfide isomerase-like enzyme. For example, treatment may include a modulation of expression of the differentially expressed genes and/or the gene encoding the human protein disulfide isomerase-like enzyme. The differential expression information may indicate whether the expression or activity of the differentially expressed gene or gene product or the
10 human protein disulfide isomerase-like gene or gene product are up-regulated or down-regulated.

Screening Methods

15 The invention provides assays for screening test compounds which bind to or modulate the activity of a protein disulfide isomerase-like polypeptide or a protein disulfide isomerase-like polynucleotide. A test compound preferably binds to a protein disulfide isomerase-like polypeptide or polynucleotide. More preferably, a test compound decreases or increases protein disulfide isomerase-like activity by at
20 least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the test compound.

Test Compounds

25 Test compounds can be pharmacologic agents already known in the art or can be compounds previously unknown to have any pharmacological activity. The compounds can be naturally occurring or designed in the laboratory. They can be isolated from microorganisms, animals, or plants, and can be produced recombinantly, or synthesized by chemical methods known in the art. If desired, test
30 compounds can be obtained using any of the numerous combinatorial library methods known in the art, including but not limited to, biological libraries, spatially

addressable parallel solid phase or solution phase libraries, synthetic library methods requiring deconvolution, the "one-bead one-compound" library method, and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer, or small molecule libraries of compounds. See Lam, *Anticancer Drug Des.* 12, 145, 1997.

Methods for the synthesis of molecular libraries are well known in the art (see, for example, DeWitt *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 90, 6909, 1993; Erb *et al.* *Proc. Natl. Acad. Sci. U.S.A.* 91, 11422, 1994; Zuckermann *et al.*, *J. Med. Chem.* 37, 2678, 1994; Cho *et al.*, *Science* 261, 1303, 1993; Carell *et al.*, *Angew. Chem. Int. Ed. Engl.* 33, 2059, 1994; Carell *et al.*, *Angew. Chem. Int. Ed. Engl.* 33, 2061; Gallop *et al.*, *J. Med. Chem.* 37, 1233, 1994). Libraries of compounds can be presented in solution (see, e.g., Houghten, *BioTechniques* 13, 412421, 1992), or on beads (Lam, *Nature* 354, 8284, 1991), chips (Fodor, *Nature* 364, 555556, 1993), bacteria or spores (Ladner, U.S. Patent 5,223,409), plasmids (Cull *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 89, 18651869, 1992), or phage (Scott & Smith, *Science* 249, 386390, 1990; Devlin, *Science* 249, 404406, 1990); Cwirla *et al.*, *Proc. Natl. Acad. Sci.* 97, 63786382, 1990; Felici, *J. Mol. Biol.* 222, 301310, 1991; and Ladner, U.S. Patent 5,223,409).

High Throughput Screening

Test compounds can be screened for the ability to bind to protein disulfide isomerase-like polypeptides or polynucleotides or to affect protein disulfide isomerase-like enzyme activity or protein disulfide isomerase-like enzyme gene expression using high throughput screening. Using high throughput screening, many discrete compounds can be tested in parallel so that large numbers of test compounds can be quickly screened. The most widely established techniques utilize 96-well microtiter plates. The wells of the microtiter plates typically require assay volumes that range from 50 to 500 μ l. In addition to the plates, many instruments, materials,

- 44 -

pipettors, robotics, plate washers, and plate readers are commercially available to fit the 96-well format.

5 Alternatively, "free format assays," or assays that have no physical barrier between samples, can be used. For example, an assay using pigment cells (melanocytes) in a simple homogeneous assay for combinatorial peptide libraries is described by Jayawickreme *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 19, 161418 (1994). The cells are placed under agarose in petri dishes, then beads that carry combinatorial compounds are placed on the surface of the agarose. The combinatorial compounds are partially released the compounds from the beads. Active compounds can be visualized as dark pigment areas because, as the compounds diffuse locally into the gel matrix, the active compounds cause the cells to change colors.

15 Another example of a free format assay is described by Chelsky, "Strategies for Screening Combinatorial Libraries: Novel and Traditional Approaches," reported at the First Annual Conference of The Society for Biomolecular Screening in Philadelphia, Pa. (Nov. 7-10, 1995). Chelsky placed a simple homogenous enzyme assay for carbonic anhydrase inside an agarose gel such that the enzyme in the gel would cause a color change throughout the gel. Thereafter, beads carrying combinatorial compounds via a photolinker were placed inside the gel and the compounds were partially released by UV light. Compounds that inhibited the enzyme were observed as local zones of inhibition having less color change.

25 Yet another example is described by Salmon *et al.*, *Molecular Diversity* 2, 5763 (1996). In this example, combinatorial libraries were screened for compounds that had cytotoxic effects on cancer cells growing in agar.

30 Another high throughput screening method is described in Beutel *et al.*, U.S. Patent 5,976,813. In this method, test samples are placed in a porous matrix. One or more assay components are then placed within, on top of, or at the bottom of a matrix such as a gel, a plastic sheet, a filter, or other form of easily manipulated solid support.

When samples are introduced to the porous matrix they diffuse sufficiently slowly, such that the assays can be performed without the test samples running together.

Binding Assays

5

For binding assays, the test compound is preferably a small molecule which binds to and occupies, for example, the ATP/GTP binding site of the enzyme or the active site of the protein disulfide isomerase-like polypeptide, such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules.

10

In binding assays, either the test compound or the protein disulfide isomerase-like polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to the protein disulfide isomerase-like polypeptide can then be accomplished, for example, by direct counting of radioemmission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product.

15

Alternatively, binding of a test compound to a protein disulfide isomerase-like polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a protein disulfide isomerase-like polypeptide. A microphysiometer (*e.g.*, Cyto-sensorJ) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and a protein disulfide isomerase-like polypeptide (McConnell *et al.*, *Science* 257, 19061912, 1992).

25

Determining the ability of a test compound to bind to a protein disulfide isomerase-like polypeptide also can be accomplished using a technology such as real-time

30

- 46 -

Bimolecular Interaction Analysis (BIA) (Sjolander & Urbaniczky, *Anal. Chem.* 63, 23382345, 1991, and Szabo *et al.*, *Curr. Opin. Struct. Biol.* 5, 699705, 1995). BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*, BIAcoreTM). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In yet another aspect of the invention, a protein disulfide isomerase-like polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent 5,283,317; Zervos *et al.*, *Cell* 72, 223232, 1993; Madura *et al.*, *J. Biol. Chem.* 268, 1204612054, 1993; Bartel *et al.*, *BioTechniques* 14, 920924, 1993; Iwabuchi *et al.*, *Oncogene* 8, 16931696, 1993; and Brent W094/10300), to identify other proteins which bind to or interact with the protein disulfide isomerase-like polypeptide and modulate its activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding a protein disulfide isomerase-like polypeptide can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to form an protein-dependent complex, the DNA-BINDING and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with the protein disulfide isomerase-like polypeptide.

It may be desirable to immobilize either the protein disulfide isomerase-like polypeptide (or polynucleotide) or the test compound to facilitate separation of bound from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the protein disulfide isomerase-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach the protein disulfide isomerase-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to a protein disulfide isomerase-like polypeptide (or polynucleotide) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, the protein disulfide isomerase-like polypeptide is a fusion protein comprising a domain that allows the protein disulfide isomerase-like polypeptide to be bound to a solid support. For example, glutathione S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the nonadsorbed protein disulfide isomerase-like polypeptide; the mixture is then incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either a protein disulfide isomerase-like polypeptide (or polynucleotide) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated protein disulfide isomerase-like polypeptides (or polynucleotides) or test compounds can be prepared from biotinNHS(Nhydroxysuccinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies which specifically bind to a protein disulfide isomerase-like polypeptide, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the ATP/GTP binding site or the active site of the protein disulfide isomerase-like polypeptide, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to the protein disulfide isomerase-like polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of the protein disulfide isomerase-like polypeptide, and SDS gel electrophoresis under non-reducing conditions.

Screening for test compounds which bind to a protein disulfide isomerase-like polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a protein disulfide isomerase-like polypeptide or polynucleotide can be used in a cell-based assay system. A protein disulfide isomerase-like polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to a protein disulfide isomerase-like polypeptide or polynucleotide is determined as described above.

Enzyme Assays

Test compounds can be tested for the ability to increase or decrease the protein disulfide isomerase-like activity of a human protein disulfide isomerase-like polypeptide. Protein disulfide isomerase-like activity can be measured, for example, as described in the specific examples, below, and in Sideraki & Gilbert, 2000, Essex *et al.*, 1995, or Chen *et al.*, 1992. Enzyme assays can be carried out after contacting either a purified protein disulfide isomerase-like polypeptide, a cell membrane preparation, or an intact cell with a test compound. A test compound which decreases a transketolase activity of a protein disulfide isomerase-like polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential therapeutic agent for decreasing protein disulfide isomerase-like enzyme activity. A test compound which increases a transketolase activity of a human protein disulfide isomerase-like polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential therapeutic agent for increasing human protein disulfide isomerase-like enzyme activity.

Gene Expression

In another embodiment, test compounds which increase or decrease protein disulfide isomerase-like enzyme gene expression are identified. A protein disulfide isomerase-like polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of the protein disulfide isomerase-like polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a modulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of

the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

5 The level of protein disulfide isomerase-like enzyme mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of a protein disulfide isomerase-like polynucleotide can be determined, for example, using a variety of techniques known
10 in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labeled amino acids into a protein disulfide isomerase-like polypeptide.

15 Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses a protein disulfide isomerase-like polynucleotide can be used in a cell-based assay system. The protein disulfide isomerase-like polynucleotide can be naturally occurring in the cell or can be introduced using
20 techniques such as those described above. Either a primary culture or an established cell line, such as CHO or human embryonic kidney 293 cells, can be used.

Pharmaceutical Compositions

25 The invention also provides pharmaceutical compositions which can be administered to a patient to achieve a therapeutic effect. Pharmaceutical compositions of the invention can comprise, for example, a protein disulfide isomerase-like polypeptide, protein disulfide isomerase-like polynucleotide, ribozymes or antisense oligonucleotides, antibodies which specifically bind to a protein disulfide isomerase-like
30 polypeptide, or mimetics, agonists, antagonists, or inhibitors of a protein disulfide isomerase-like polypeptide activity. The compositions can be administered alone or

- 51 -

in combination with at least one other agent, such as stabilizing compound, which can be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions can be administered to a patient alone, or in combination with other agents, drugs or hormones.

In addition to the active ingredients, these pharmaceutical compositions can contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Pharmaceutical compositions of the invention can be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intraarterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, parenteral, topical, sublingual, or rectal means. Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethylcellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

30

- 52 -

Dragee cores can be used in conjunction with suitable coatings, such as concentrated sugar solutions, which also can contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, *i.e.*, dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers also can be used for delivery. Optionally, the suspension also can contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

- 53 -

The pharmaceutical compositions of the present invention can be manufactured in a manner that is known in the art, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. The pharmaceutical composition can be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation can be a lyophilized powder which can contain any or all of the following: 150 mM histidine, 0.1%2% sucrose, and 27% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

Further details on techniques for formulation and administration can be found in the latest edition of REMINGTON'S PHARMACEUTICAL SCIENCES (Maack Publishing Co., Easton, Pa.). After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. Such labeling would include amount, frequency, and method of administration.

Therapeutic Indications

PDI has been associated with failure of potent anti-proliferative proteins such as TGF β 1 to inhibit proliferation of cancer cells. It is postulated that PDI acts by altering the binding of either the protein or its receptor. A therapeutic strategy targeted toward reducing the expression or activity of human protein disulfide isomerase-like enzyme, therefore, could enhance the anti-proliferative effective of anti-neoplastic compounds, particularly compounds which require specific protein conformations for maximum anti-proliferative effects for the treatment of cancer.

Cancer is a disease fundamentally caused by oncogenic cellular transformation. There are several hallmarks of transformed cells that distinguish them from their normal counterparts and underlie the pathophysiology of cancer. These include

- 54 -

uncontrolled cellular proliferation, unresponsiveness to normal death inducing signals (immortalization), increased cellular motility and invasiveness, increased ability to recruit blood supply through induction of new blood vessel formation (angiogenesis), genetic instability, and dysregulated gene expression. Various combinations of these aberrant physiologies, along with the acquisition of drug-resistance frequently lead to an intractable disease state in which organ failure and patient death ultimately ensue.

The advent of genomics-driven molecular target identification has opened up the possibility of identifying new cancer-specific targets for therapeutic intervention that will provide safer, more effective treatments for cancer patients. Thus, newly discovered tumor-associated genes and their products can be tested for their role(s) in disease and used as tools to discover and develop innovative therapies.

Genes playing important roles in any of the physiological processes outlined above can be characterized as cancer targets. Genes or gene fragments identified through genomics can readily be expressed in one or more heterologous expression systems to produce functional recombinant proteins. These proteins are characterized *in vitro* for their biochemical properties and then used as tools in high-throughput molecular screening programs to identify chemical modulators of their biochemical activities. Agonists and/or antagonists of target protein activity can be identified in this manner and subsequently tested in cellular and *in vivo* disease models for anticancer activity. Optimization of lead compounds with iterative testing in biological models and detailed pharmacokinetic and toxicological analyses form the basis for drug development and subsequent testing in humans.

Human protein disulfide isomerase-like enzyme also can be regulated to treat hematological disorders and atherosclerosis. Atherosclerosis is associated with abnormally high level of PDI. Therefore, blocking the activity of human protein disulfide isomerase-like enzyme would be a viable approach to treating or preventing atherosclerosis.

Human protein disulfide isomerase-like enzyme also can be regulated to treat other cardiovascular disorders. Cardiovascular diseases include the following disorders of the heart and the vascular system: congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases and peripheral vascular diseases.

Heart failure is defined as a pathophysiologic state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failure such as high-output and low-output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

Ischemic diseases are conditions in which the coronary flow is restricted resulting in an perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases include stable angina, unstable angina and asymptomatic ischemia.

Arrhythmias include all forms of atrial and ventricular tachyarrhythmias (atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation) as well as bradycardic forms of arrhythmias.

Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hypertension (renal, endocrine, neurogenic, others). The genes may be used as drug targets for the treatment of hypertension as well as for the prevention of all

5 complications. Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

10 PDI is postulated to play a role in wound healing, because PDI is released from activated platelets. PDI may promote tissue remodeling and arrest of bleeding by reduction or isomerization of protein bonds. By enhancing the expression of human protein disulfide isomerase-like enzyme, it may be possible to increase the levels of properly folded proteins that are important in facilitating wound healing.

15 This invention further pertains to the use of novel agents identified by the screening assays described above. Accordingly, it is within the scope of this invention to use a test compound identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a modulating agent, an antisense nucleic acid molecule, a specific antibody, ribozyme, or a human protein disulfide isomerase-like polypeptide binding molecule) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above described screening assays for treatments as described herein.

25 A reagent which affects human protein disulfide isomerase-like enzyme activity can be administered to a human cell, either *in vitro* or *in vivo*, to reduce human protein disulfide isomerase-like enzyme activity. The reagent preferably binds to an expression product of a human protein disulfide isomerase-like enzyme gene. If the expression product is a protein, the reagent is preferably an antibody. For treatment of human cells *ex vivo*, an antibody can be added to a preparation of stem cells which

30

- 57 -

have been removed from the body. The cells can then be replaced in the same or another human body, with or without clonal propagation, as is known in the art.

5 In one embodiment, the reagent is delivered using a liposome. Preferably, the liposome is stable in the animal into which it has been administered for at least about 30 minutes, more preferably for at least about 1 hour, and even more preferably for at least about 24 hours. A liposome comprises a lipid composition that is capable of targeting a reagent, particularly a polynucleotide, to a particular site in an animal, such as a human. Preferably, the lipid composition of the liposome is capable of
10 targeting to a specific organ of an animal, such as the lung, liver, spleen, heart brain, lymph nodes, and skin.

A liposome useful in the present invention comprises a lipid composition that is capable of fusing with the plasma membrane of the targeted cell to deliver its
15 contents to the cell. Preferably, the transfection efficiency of a liposome is about 0.5 μg of DNA per 16 nmole of liposome delivered to about 10^6 cells, more preferably about 1.0 μg of DNA per 16 nmole of liposome delivered to about 10^6 cells, and even more preferably about 2.0 μg of DNA per 16 nmol of liposome delivered to about 10^6 cells. Preferably, a liposome is between about 100 and
20 500 nm, more preferably between about 150 and 450 nm, and even more preferably between about 200 and 400 nm in diameter.

Suitable liposomes for use in the present invention include those liposomes standardly used in, for example, gene delivery methods known to those of skill in the
25 art. More preferred liposomes include liposomes having a polycationic lipid composition and/or liposomes having a cholesterol backbone conjugated to polyethylene glycol. Optionally, a liposome comprises a compound capable of targeting the liposome to a particular cell type, such as a cell-specific ligand exposed on the outer surface of the liposome.

30

Complexing a liposome with a reagent such as an antisense oligonucleotide or ribozyme can be achieved using methods which are standard in the art (see, for example, U.S. Patent 5,705,151). Preferably, from about 0.1 μ g to about 10 μ g of polynucleotide is combined with about 8 nmol of liposomes, more preferably from about 0.5 μ g to about 5 μ g of polynucleotides are combined with about 8 nmol liposomes, and even more preferably about 1.0 μ g of polynucleotides is combined with about 8 nmol liposomes.

In another embodiment, antibodies can be delivered to specific tissues *in vivo* using receptor-mediated targeted delivery. Receptor-mediated DNA delivery techniques are taught in, for example, Findeis *et al.* *Trends in Biotechnol.* 11, 202-05 (1993); Chiou *et al.*, GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER (J.A. Wolff, ed.) (1994); Wu & Wu, *J. Biol. Chem.* 263, 621-24 (1988); Wu *et al.*, *J. Biol. Chem.* 269, 542-46 (1994); Zenke *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 87, 3655-59 (1990); Wu *et al.*, *J. Biol. Chem.* 266, 338-42 (1991).

Determination of a Therapeutically Effective Dose

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases human protein disulfide isomerase-like enzyme activity relative to the human protein disulfide isomerase-like enzyme activity which occurs in the absence of the therapeutically effective dose.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, *e.g.*, ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it
5 can be expressed as the ratio, LD₅₀/ED₅₀.

Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is
10 preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to
15 the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and
20 tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 to 100,000 micrograms, up to a total dose
25 of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations,
30 etc.

- 60 -

If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun," and DEAE- or calcium phosphate-mediated transfection.

Effective *in vivo* dosages of an antibody are in the range of about 5 µg to about 50 µg/kg, about 50 µg to about 5 mg/kg, about 100 µg to about 500 µg/kg of patient body weight, and about 200 to about 250 µg/kg of patient body weight. For administration of polynucleotides encoding single-chain antibodies, effective *in vivo* dosages are in the range of about 100 ng to about 200 ng, 500 ng to about 50 mg, about 1 µg to about 2 mg, about 5 µg to about 500 µg, and about 20 µg to about 100 µg of DNA.

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above.

Preferably, a reagent reduces expression of a human protein disulfide isomerase-like enzyme gene or the activity of a human protein disulfide isomerase-like polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of a human protein disulfide isomerase-like enzyme gene or the activity of a human protein disulfide isomerase-like polypeptide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to human protein disulfide isomerase-like enzyme-specific mRNA, quantitative RT-PCR, immunologic detection of a human protein disulfide

- 61 -

isomerase-like polypeptide, or measurement of human protein disulfide isomerase-like enzyme activity.

5 In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described
10 above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows,
15 horses, rabbits, monkeys, and most preferably, humans.

Diagnostic Methods

Human protein disulfide isomerase-like enzyme also can be used in diagnostic assays
20 for detecting diseases and abnormalities or susceptibility to diseases and abnormalities related to the presence of mutations in the nucleic acid sequences which encode the enzyme. For example, differences can be determined between the cDNA or genomic sequence encoding protein disulfide isomerase-like enzyme in individuals afflicted with a disease and in normal individuals. If a mutation is
25 observed in some or all of the afflicted individuals but not in normal individuals, then the mutation is likely to be the causative agent of the disease.

Sequence differences between a reference gene and a gene having mutations can be revealed by the direct DNA sequencing method. In addition, cloned DNA segments
30 can be employed as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with PCR. For example, a sequencing

- 62 -

primer can be used with a double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures using radiolabeled nucleotides or by automatic sequencing procedures using fluorescent tags.

5

Genetic testing based on DNA sequence differences can be carried out by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized, for example, by high resolution gel electrophoresis. DNA fragments of different sequences can be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (*see, e.g., Myers et al., Science 230, 1242, 1985*). Sequence changes at specific locations can also be revealed by nuclease protection assays, such as RNase and S 1 protection or the chemical cleavage method (*e.g., Cotton et al., Proc. Natl. Acad. Sci. USA 85, 43974401, 1985*). Thus, the detection of a specific DNA sequence can be performed by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes and Southern blotting of genomic DNA. In addition to direct methods such as gel electrophoresis and DNA sequencing, mutations can also be detected by *in situ* analysis.

20

Altered levels of a protein disulfide isomerase-like enzyme also can be detected in various tissues. Assays used to detect levels of the receptor polypeptides in a body sample, such as blood or a tissue biopsy, derived from a host are well known to those of skill in the art and include radioimmunoassays, competitive binding assays, Western blot analysis, and ELISA assays.

25

All patents and patent applications cited in this disclosure are expressly incorporated herein by reference. The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following

30

specific examples which are provided for purposes of illustration only and are not intended to limit the scope of the invention.

EXAMPLE 1

Detection of protein disulfide isomerase-like enzyme activity

The polynucleotide of SEQ ID NO:1 or SEQ ID NO:28 is inserted into the expression vector pCEV4 and the expression vector pCEV4-protein disulfide isomerase-like enzyme polypeptide obtained is transfected into human embryonic kidney 293 cells. From these cells extracts are obtained and the protein disulfide isomerase-like enzyme activity is measured by it's ability to refold scrambled Rnase A, which in turn will regain its ability to cleave highly polymerized yeast RNA (Lambert and Freedman, 1983). The cell extract (2,0µg/ml) is dissolved in 100 mM sodium phosphate, pH 7,5, 2 mM glutathione, and 0,2 mM reduced glutathione. The reaction is started by the addition of 200 µg of scrambled Rnase A (Hillson et al., 1984). Protein disulfide isomerase-like enzyme activity is determined in the presence of recombinant calretioulin, purified N-, P-, and C-domains of calreticulin, and in the presence of cardiac calsequestrin. For the assay, the proteins are used in 1:1 molar ratio. The A260/A280 ratio over a period of 60 min is measured as indicative of the extent of cleavage of yeast RNA by refolded Rnase A. It is shown that the polypeptide of SEQ ID NO:2 and SEQ ID NO:29 has a protein disulfide isomerase-like enzyme activity.

EXAMPLE 2

Expression of recombinant human protein disulfide isomerase-like enzyme

The *Pichia pastoris* expression vector pPICZB (Invitrogen, San Diego, CA) is used to produce large quantities of recombinant human protein disulfide isomerase-like polypeptides in yeast. The protein disulfide isomerase-like enzyme-encoding DNA

- 64 -

sequence is derived from SEQ ID NO:1 or SEQ ID NO:28. Before insertion into vector pPICZB, the DNA sequence is modified by well known methods in such a way that it contains at its 5' end an initiation codon and at its 3' end an enterokinase cleavage site, a His6 reporter tag and a termination codon. Moreover, at both termini
5 recognition sequences for restriction endonucleases are added and after digestion of the multiple cloning site of pPICZ B with the corresponding restriction enzymes the modified DNA sequence is ligated into pPICZB. This expression vector is designed for inducible expression in *Pichia pastoris*, driven by a yeast promoter. The resulting pPICZ/mdHis6 vector is used to transform the yeast

10 The yeast is cultivated under usual conditions in 5 liter shake flasks and the recombinantly produced protein isolated from the culture by affinity chromatography (NiNTA Resin) in the presence of 8 M urea. The bound polypeptide is eluted with buffer, pH 3.5, and neutralized. Separation of the polypeptide from the His6 reporter
15 tag is accomplished by site-specific proteolysis using enterokinase (Invitrogen, San Diego, CA) according to manufacturer's instructions. Purified human protein disulfide isomerase-like polypeptide is obtained.

EXAMPLE 3

20 *Identification of a test compound which decreases protein disulfide isomerase-like activity*

Human protein disulfide isomerase-like enzyme activity is measured by renaturation
25 of inactivated RNase. Briefly, RNase is inactivated by formation of randomly mismatched or scrambled disulfide bonds. Deactivated RNase renatures slowly in the presence of a thiol reagent. Human protein disulfide isomerase-like enzyme accelerates the renaturation process. The amount of renatured RNase is measured by its ability to digest RNA. RNA digestion is measured by an increase in absorbance at
30 260 nm.

- 65 -

A test compound which decreases protein disulfide isomerase-like activity relative to control levels by at least 20% is identified as a protein disulfide isomerase-like enzyme inhibitor.

5 **EXAMPLE 4**

Identification of test compounds that bind to protein disulfide isomerase-like polypeptides

10 Purified protein disulfide isomerase-like polypeptides comprising a glutathione-S-transferase protein and absorbed onto glutathione-derivatized wells of 96-well microtiter plates are contacted with test compounds from a small molecule library at pH 7.0 in a physiological buffer solution. Protein disulfide isomerase-like polypeptides comprise the amino acid sequence shown in SEQ ID NO:2 or SEQ ID
15 NO:29. The test compounds comprise a fluorescent tag. The samples are incubated for 5 minutes to one hour. Control samples are incubated in the absence of a test compound.

The buffer solution containing the test compounds is washed from the wells.
20 Binding of a test compound to a protein disulfide isomerase-like polypeptide is detected by fluorescence measurements of the contents of the wells. A test compound which increases the fluorescence in a well by at least 15% relative to fluorescence of a well in which a test compound is not incubated is identified as a compound which binds to a protein disulfide isomerase-like polypeptide.

25

EXAMPLE 5

Identification of a test compound which decreases protein disulfide isomerase-like enzyme gene expression

30

- 66 -

5 A test compound is administered to a culture of human cells transfected with a protein disulfide isomerase-like enzyme expression construct and incubated at 37°C for 10 to 45 minutes. A culture of the same type of cells which have not been transfected is incubated for the same time without the test compound to provide a negative control.

10 RNA is isolated from the two cultures as described in Chirgwin *et al.*, *Biochem. 18*, 5294-99, 1979). Northern blots are prepared using 20 to 30 Fg total RNA and hybridized with a ³²P-labeled protein disulfide isomerase-like enzyme-specific probe at 65° C in Express-hyb (CLONTECH). The probe comprises at least 11 contiguous nucleotides selected from the complement of SEQ ID NO:1 or SEQ ID NO:28. A test compound which decreases the protein disulfide isomerase-like enzyme-specific signal relative to the signal obtained in the absence of the test compound is identified as an inhibitor of protein disulfide isomerase-like enzyme gene expression.

15

EXAMPLE 6

Proliferation inhibition assay: Antisense oligonucleotides suppress the growth of cancer cell lines in the presence of TGFβ1

20

A human pancreatic carcinoma line which does not downregulate the expression of human protein disulfide isomerase-like enzyme in response to TGFβ1 and which is resistant to the anti-proliferative effects of TGFβ1 is used for testing. Cells are cultured in RPMI-1640 with 10-15% fetal calf serum at a concentration of 10,000 cells per milliliter in a volume of 0.5 ml and kept at 37 °C in a 95% air/5%CO₂ atmosphere.

25

30 Phosphorothioate oligoribonucleotides are synthesized on an Applied Biosystems Model 380B DNA synthesizer using phosphoroamidite chemistry. A sequence of 24 bases complementary to the nucleotides at position 1 to 24 of SEQ ID NO:1 or SEQ ID NO:28 is used as the test oligonucleotide. As a control, another (random)

- 67 -

sequence is used: 5'-TCA ACT GAC TAG ATG TAC ATG GAC-3'. Following assembly and deprotection, oligonucleotides are ethanol-precipitated twice, dried, and suspended in phosphate buffered saline at the desired concentration. Purity of the oligonucleotides is tested by capillary gel electrophoresis and ion exchange HPLC. The purified oligonucleotides are added to the culture medium at a concentration of 10 μ M once per day for seven days.

The addition of the test oligonucleotide for seven days together with TGF β 1 results in significantly reduced expression of human protein disulfide isomerase-like enzyme as determined by Western blotting. This effect is not observed with the control oligonucleotide. After 3 to 7 days, the number of cells in the cultures is counted using an automatic cell counter. The number of cells in cultures treated with the test oligonucleotide (expressed as 100%) is compared with the number of cells in cultures treated with the control oligonucleotide. The number of cells in cultures treated with the test oligonucleotide is not more than 30% of control, indicating that the inhibition of human protein disulfide isomerase-like enzyme has an anti-proliferative effect on cancer cells.

EXAMPLE 7

Quantitative expression profiling of human protein disulfide isomerase-like enzyme

Expression profiling is based on a quantitative polymerase chain reaction (PCR) analysis, also called kinetic analysis, first described in Higuchi et al., 1992 and Higuchi et al., 1993. The principle is that at any given cycle within the exponential phase of PCR, the amount of product is proportional to the initial number of template copies. Using this technique, the expression levels of particular genes, which are transcribed from the chromosomes as messenger RNA (mRNA), are measured by first making a DNA copy (cDNA) of the mRNA, and then performing quantitative PCR on the cDNA, a method called quantitative reverse transcription-polymerase chain reaction (quantitative RT-PCR).

- 68 -

Quantitative RT-PCR analysis of RNA from different human tissues was performed to investigate the tissue distribution of human protein disulfide isomerase-like enzyme, Sphingosine kinase-like mRNA. In most cases, 25 µg of total RNA from various tissues (including Human Total RNA Panel I-V, Clontech Laboratories, Palo Alto, CA, USA) was used as a template to synthesize first-strand cDNA using the SUPERScript™ First-Strand Synthesis System for RT-PCR (Life Technologies, Rockville, MD, USA). First-strand cDNA synthesis was carried out according to the manufacturer's protocol using oligo (dT) to hybridize to the 3' poly A tails of mRNA and prime the synthesis reaction. Approximately 10 ng of the first-strand cDNA was then used as template in a polymerase chain reaction. In other cases, 10 ng of commercially available cDNAs (Human Immune System MTC Panel and Human Blood Fractions MTC Panel, Clontech Laboratories, Palo Alto, CA, USA) were used as template in a polymerase chain reaction. The polymerase chain reaction was performed in a LightCycler (Roche Molecular Biochemicals, Indianapolis, IN, USA), in the presence of the DNA-binding fluorescent dye SYBR Green I which binds to the minor groove of the DNA double helix, produced only when double-stranded DNA is successfully synthesized in the reaction. Upon binding to double-stranded DNA, SYBR Green I emits light that can be quantitatively measured by the LightCycler machine. The polymerase chain reaction was carried out using oligonucleotide primers LBRI059-L2:

GGACACATTCACCTGTGGCAAGG

and LBRI059-R2:

TGGAAGCAGGG TCTGAGTGATG

and measurements of the intensity of emitted light were taken following each cycle of the reaction when the reaction had reached a temperature of 81 degrees C. Intensities of emitted light were converted into copy numbers of the gene transcript per nanogram of template cDNA by comparison with simultaneously reacted standards of known concentration.

To correct for differences in mRNA transcription levels per cell in the various tissue types, a normalization procedure was performed using similarly calculated expression levels in the various tissues of five different housekeeping genes: glycerinaldehyde-3-phosphatase (G3PDH), hypoxanthine guanine phosphoribosyl transferase (HPRT), beta-actin, porphobilinogen deaminase (PBGD), and beta-2-microglobulin. The level of housekeeping gene expression is considered to be relatively constant for all tissues and therefore can be used as a gauge to approximate relative numbers of cells per μg of total RNA used in the cDNA synthesis step. Except for the use of a slightly different set of housekeeping genes and the use of the LightCycler system to measure expression levels, the normalization procedure was similar to that described in the RNA Master Blot User Manual, Appendix C (1997, Clontech Laboratories, Palo Alto, CA, USA). In brief, expression levels of the five housekeeping genes in all tissue samples were measured in three independent reactions per gene using the LightCycler and a constant amount (25 μg) of starting RNA. The calculated copy numbers for each gene, derived from comparison with simultaneously reacted standards of known concentrations, were recorded and the mean number of copies of each gene in all tissue samples was determined. Then for each tissue sample, the expression of each housekeeping gene relative to the mean was calculated, and the average of these values over the five housekeeping genes was found. A normalization factor for each tissue was then calculated by dividing the final value for one of the tissues arbitrarily selected as a standard by the corresponding value for each of the tissues. To normalize an experimentally obtained value for the expression of a particular gene in a tissue sample, the obtained value was multiplied by the normalization factor for the tissue tested. This normalization method was used for all tissues except those derived from the Human Blood Fractions MTC Panel, which showed dramatic variation in some housekeeping genes depending on whether the tissue had been activated or not. In these tissues, normalization was carried out with a single housekeeping gene, beta-2-microglobulin.

- 70 -

Results are given in Figures 35 and 36, showing the experimentally obtained copy numbers of mRNA per 10 ng of first-strand cDNA on the left and the normalized values on the right. RNAs used for the cDNA synthesis, along with their supplier and catalog numbers are shown in tables 1 and 2.

5

Human protein disulfide isomerase-like enzyme is a widely distributed, highly expressed molecule, indicating that its function may be fundamental to common activities of a wide variety of cell types. As a member of the protein disulfide isomerase family, this function would most likely be in helping different proteins

10 assume their proper three-dimensional configuration.

15

Although the expression of human protein disulfide isomerase-like enzyme is high in most tissues, its expression in the lung is 1.5 to 2 times that seen in any other tissue. This suggests that the lung may produce a comparatively large amount of specific protein types, for example proteins lining or secreted into the lung airways, that require human protein disulfide isomerase-like enzyme activity to allow them to fold or combine with other molecules before assuming an active conformation. Regulating the activity of human protein disulfide isomerase-like enzyme in cells or tissues such as lung that produce human protein disulfide isomerase-like enzyme -

20 activity-dependent proteins, therefore, may be beneficial in pathogenic states where overproduction or underproduction of active protein leads to disease.

Table 1: Material used to determine the expression in whole body tissues.

Tissue	Supplier	Panel name and catalog number
1. brain	Clontech	Human Total RNA Panel I, K4000-1
2. heart	Clontech	Human Total RNA Panel I, K4000-1
3. kidney	Clontech	Human Total RNA Panel I, K4000-1
4. liver	Clontech	Human Total RNA Panel I, K4000-1
5. lung	Clontech	Human Total RNA Panel I, K4000-1
6. trachea	Clontech	Human Total RNA Panel I, K4000-1
7. bone marrow	Clontech	Human Total RNA Panel II, K4001-1
8. colon	Clontech	Human Total RNA Panel II, K4001-1
9. small intestine	Clontech	Human Total RNA Panel II, K4001-1
10. spleen	Clontech	Human Total RNA Panel II, K4001-1
11. stomach	Clontech	Human Total RNA Panel II, K4001-1
12. thymus	Clontech	Human Total RNA Panel II, K4001-1
13. mammary gland	Clontech	Human Total RNA Panel III, K4002-1
14. skeletal muscle	Clontech	Human Total RNA Panel III, K4002-1
15. prostate	Clontech	Human Total RNA Panel III, K4002-1
16. testis	Clontech	Human Total RNA Panel III, K4002-1
17. uterus	Clontech	Human Total RNA Panel III, K4002-1
18. cerebellum	Clontech	Human Total RNA Panel IV, K4003-1
19. fetal brain	Clontech	Human Total RNA Panel IV, K4003-1
20. fetal liver	Clontech	Human Total RNA Panel IV, K4003-1
21. spinal cord	Clontech	Human Total RNA Panel IV, K4003-1
22. placenta	Clontech	Human Total RNA Panel IV, K4003-1
23. adrenal gland	Clontech	Human Total RNA Panel V, K4004-1
24. pancreas	Clontech	Human Total RNA Panel V, K4004-1
25. salivary gland	Clontech	Human Total RNA Panel V, K4004-1
26. thyroid	Clontech	Human Total RNA Panel V, K4004-1

Table 2: Material used to determine the expression in the blood and lung tissues.

Tissue	Supplier	Panel name and catalog number
1. lymph node	Clontech	Human Immune System MTC Panel, K1426-1
2. peripheral blood leukocytes	Clontech	Human Immune System MTC Panel, K1426-1
3. tonsil	Clontech	Human Immune System MTC Panel, K1426-1
4. peripheral blood mononuclear cells	Clontech	Human Blood Fractions MTC Panel, K1428-1
5. peripheral blood mononuclear cells – activated	Clontech	Human Blood Fractions MTC Panel, K1428-1
6. T-cell (CD8+)	Clontech	Human Blood Fractions MTC Panel, K1428-1
7. T-cell (CD8+) - activated	Clontech	Human Blood Fractions MTC Panel, K1428-1
8. T-cell (CD4+)	Clontech	Human Blood Fractions MTC Panel, K1428-1
9. T-cell (CD4+) – activated	Clontech	Human Blood Fractions MTC Panel, K1428-1
10. B-cell (CD19+)	Clontech	Human Blood Fractions MTC Panel, K1428-1
11. B-cell (CD19+) – activated	Clontech	Human Blood Fractions MTC Panel, K1428-1
12. Monocytes (CD14+)	Clontech	Human Blood Fractions MTC Panel, K1428-1
13. Th1 clone	In-house	
14. Th2 clone	In-house	
15. neutrophil	In-house	
16. neutrophil	In-house	
17. Normal Bronchial/Tracheal Epithelial Cells	In-house	
18. Normal Bronchial/Tracheal smooth muscle cell	In-house	
19. Normal lung fibroblast	In-house	
20. Microvascular Endothelial cell	In-house	
21. U937	In-house	
22. RAMOS	In-house	
23. Jurkat	In-house	
24. HeLaS3	In-house	
25. IMR-90	In-house	
26. HEK293	In-house	

CLAIMS

1. An isolated polynucleotide encoding a protein disulfide isomerase-like enzyme polypeptide and being selected from the group consisting of:
- 5
- a) a polynucleotide encoding a protein disulfide isomerase-like enzyme polypeptide comprising an amino acid sequence selected from the group consisting of:
- 10 amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:2, amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:29, the amino acid sequence shown in SEQ ID NO:2; and
- 15 the amino acid sequence shown in SEQ ID NO:29.
- b) a polynucleotide comprising the sequence of SEQ ID NO:1 or SEQ ID NO:28;
- 20 c) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) and (b);
- d) a polynucleotide the sequence of which deviates from the polynucleotide sequences specified in (a) to (c) due to the degeneration of
- 25 the genetic code; and
- e) a polynucleotide which represents a fragment, derivative or allelic variation of a polynucleotide sequence specified in (a to (d).
- 30 2. An expression vector containing any polynucleotide of claim 1.

3. A host cell containing the expression vector of claim 2.
4. A substantially purified protein disulfide isomerase-like enzyme polypeptide encoded by a polynucleotide of claim 1.
5. A method for producing a protein disulfide isomerase-like enzyme polypeptide, wherein the method comprises the following steps:
 - 10 a) culturing the host cell of claim 3 under conditions suitable for the expression of the protein disulfide isomerase-like enzyme polypeptide; and
 - b) recovering the protein disulfide isomerase-like enzyme polypeptide from the host cell culture.
- 15 6. A method for detection of a polynucleotide encoding a protein disulfide isomerase-like enzyme polypeptide in a biological sample comprising the following steps:
 - 20 a) hybridizing any polynucleotide of claim 1 to a nucleic acid material of a biological sample, thereby forming a hybridization complex; and
 - b) detecting said hybridization complex.
- 25 7. The method of claim 6, wherein before hybridization, the nucleic acid material of the biological sample is amplified.
8. A method for the detection of a polynucleotide of claim 1 or a protein disulfide isomerase-like enzyme polypeptide of claim 4 comprising the steps of:

- 75 -

contacting a biological sample with a reagent which specifically interacts with the polynucleotide or the protein disulfide isomerase-like enzyme polypeptide.

- 5 9. A diagnostic kit for conducting the method of any one of claims 6 to 8.
10. A method of screening for agents which decrease the activity of a protein disulfide isomerase-like enzyme, comprising the steps of:
- 10 contacting a test compound with any protein disulfide isomerase-like enzyme polypeptide encoded by any polynucleotide of claim 1;
- 15 detecting binding of the test compound to the protein disulfide isomerase-like enzyme polypeptide, wherein a test compound which binds to the polypeptide is identified as a potential therapeutic agent for decreasing the activity of a protein disulfide isomerase-like enzyme.
11. A method of screening for agents which regulate the activity of a protein disulfide isomerase-like enzyme, comprising the steps of:
- 20 contacting a test compound with a protein disulfide isomerase-like enzyme polypeptide encoded by any polynucleotide of claim 1; and
- 25 detecting a protein disulfide isomerase-like enzyme activity of the polypeptide, wherein a test compound which increases the protein disulfide isomerase-like enzyme activity is identified as a potential therapeutic agent for increasing the activity of the protein disulfide isomerase-like enzyme, and wherein a test compound which decreases the protein disulfide isomerase-like enzyme activity of the polypeptide is identified as a potential therapeutic agent for decreasing the activity of the protein disulfide isomerase-like enzyme.
- 30

12. A method of screening for agents which decrease the activity of a protein disulfide isomerase-like enzyme, comprising the steps of:

5 contacting a test compound with any polynucleotide of claim 1 and detecting binding of the test compound to the polynucleotide, wherein a test compound which binds to the polynucleotide is identified as a potential therapeutic agent for decreasing the activity of protein disulfide isomerase-like enzyme.

- 10 13. A method of reducing the activity of protein disulfide isomerase-like enzyme, comprising the steps of:

15 contacting a cell with a reagent which specifically binds to any polynucleotide of claim 1 or any protein disulfide isomerase-like enzyme polypeptide of claim 4, whereby the activity of protein disulfide isomerase-like enzyme is reduced.

- 20 14. A reagent that modulates the activity of a protein disulfide isomerase-like enzyme polypeptide or a polynucleotide wherein said reagent is identified by the method of any of the claim 10 to 12.

15. A pharmaceutical composition, comprising:

25 the expression vector of claim 2 or the reagent of claim 14 and a pharmaceutically acceptable carrier.

16. Use of the pharmaceutical composition of claim 15 for modulating the activity of a protein disulfide isomerase-like enzyme in a disease.

- 30 17. Use of claim 16 wherein the disease is cancer, diabetes and obesity.

- 77 -

18. A cDNA encoding a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29.
19. The cDNA of claim 18 which comprises SEQ ID NO:1 or SEQ ID NO:28.
20. The cDNA of claim 18 which consists of SEQ ID NO:1 or SEQ ID NO:28.
21. An expression vector comprising a polynucleotide which encodes a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29.
22. The expression vector of claim 21 wherein the polynucleotide consists of SEQ ID NO:1 or SEQ ID NO:28.
23. A host cell comprising an expression vector which encodes a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29.
24. The host cell of claim 23 wherein the polynucleotide consists of SEQ ID NO:1 or SEQ ID NO:28.
25. A purified polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29.
26. The purified polypeptide of claim 25 which consists of the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29.
27. A fusion protein comprising a polypeptide having the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29.

- 78 -

28. A method of producing a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29, comprising the steps of:

5 culturing a host cell comprising an expression vector which encodes the polypeptide under conditions whereby the polypeptide is expressed; and

isolating the polypeptide.

29. The method of claim 28 wherein the expression vector comprises SEQ ID
10 NO:1 or SEQ ID NO:28.

30. A method of detecting a coding sequence for a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29, comprising the steps of:

15 hybridizing a polynucleotide comprising 11 contiguous nucleotides of SEQ ID NO:1 or SEQ ID NO:28 to nucleic acid material of a biological sample, thereby forming a hybridization complex; and

20 detecting the hybridization complex.

31. The method of claim 30 further comprising the step of amplifying the nucleic acid material before the step of hybridizing.

- 25 32. A kit for detecting a coding sequence for a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29, comprising:

a polynucleotide comprising 11 contiguous nucleotides of SEQ ID NO:1 or
30 SEQ ID NO:28; and

instructions for the method of claim 30.

33. A method of detecting a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29, comprising the steps of:
- 5 contacting a biological sample with a reagent that specifically binds to the polypeptide to form a reagent-polypeptide complex; and
- detecting the reagent-polypeptide complex.
- 10 34. The method of claim 33 wherein the reagent is an antibody.
35. A kit for detecting a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29, comprising:
- 15 an antibody which specifically binds to the polypeptide; and
- instructions for the method of claim 33.
- 20 36. A method of screening for agents which can modulate the activity of a human protein disulfide isomerase-like enzyme, comprising the steps of:
- contacting a test compound with a polypeptide comprising an amino acid sequence selected from the group consisting of: (1) amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:2 or which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:29 and (2) the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29; and
- 25 detecting binding of the test compound to the polypeptide, wherein a test compound which binds to the polypeptide is identified as a potential agent for regulating activity of the human protein disulfide isomerase-like enzyme.
- 30

37. The method of claim 36 wherein the step of contacting is in a cell.
38. The method of claim 36 wherein the cell is *in vitro*.
- 5 39. The method of claim 36 wherein the step of contacting is in a cell-free system.
40. The method of claim 36 wherein the polypeptide comprises a detectable label.
- 10 41. The method of claim 36 wherein the test compound comprises a detectable label.
42. The method of claim 36 wherein the test compound displaces a labeled ligand which is bound to the polypeptide.
- 15 43. The method of claim 36 wherein the polypeptide is bound to a solid support.
44. The method of claim 36 wherein the test compound is bound to a solid support.
- 20 45. A method of screening for agents which modulate an activity of a human protein disulfide isomerase-like enzyme, comprising the steps of:
- 25 contacting a test compound with a polypeptide comprising an amino acid sequence selected from the group consisting of: (1) amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:2 or which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO:29 and (2) the amino acid sequence shown in
- 30 SEQ ID NO:2 or in SEQ ID NO:29; and

detecting an activity of the polypeptide, wherein a test compound which increases the activity of the polypeptide is identified as a potential agent for increasing the activity of the human protein disulfide isomerase-like enzyme, and wherein a test compound which decreases the activity of the polypeptide is identified as a potential agent for decreasing the activity of the human protein disulfide isomerase-like enzyme.

- 5
46. The method of claim 45 wherein the step of contacting is in a cell.
- 10
47. The method of claim 45 wherein the cell is *in vitro*.
48. The method of claim 45 wherein the step of contacting is in a cell-free system.
- 15
49. A method of screening for agents which modulate an activity of a human protein disulfide isomerase-like enzyme, comprising the steps of:
- 20
- contacting a test compound with a product encoded by a polynucleotide which comprises the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:28; and
- 25
- detecting binding of the test compound to the product, wherein a test compound which binds to the product is identified as a potential agent for regulating the activity of the human protein disulfide isomerase-like enzyme.
50. The method of claim 49 wherein the product is a polypeptide.
51. The method of claim 49 wherein the product is RNA.
- 30
52. A method of reducing activity of a human protein disulfide isomerase-like enzyme, comprising the step of:

contacting a cell with a reagent which specifically binds to a product encoded by a polynucleotide comprising the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:28, whereby the activity of a human protein disulfide isomerase-like enzyme is reduced.

53. The method of claim 52 wherein the product is a polypeptide.

54. The method of claim 53 wherein the reagent is an antibody.

55. The method of claim 52 wherein the product is RNA.

56. The method of claim 55 wherein the reagent is an antisense oligonucleotide.

57. The method of claim 56 wherein the reagent is a ribozyme.

58. The method of claim 52 wherein the cell is *in vitro*.

59. The method of claim 52 wherein the cell is *in vivo*.

60. A pharmaceutical composition, comprising:

a reagent which specifically binds to a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29; and

a pharmaceutically acceptable carrier.

61. The pharmaceutical composition of claim 60 wherein the reagent is an antibody.

62. A pharmaceutical composition, comprising:

a reagent which specifically binds to a product of a polynucleotide comprising the nucleotide sequence shown in SEQ ID NO:1 or SEQ ID NO:28; and a pharmaceutically acceptable carrier.

5 63. The pharmaceutical composition of claim 62 wherein the reagent is a ribozyme.

64. The pharmaceutical composition of claim 62 wherein the reagent is an antisense oligonucleotide.

10

65. The pharmaceutical composition of claim 62 wherein the reagent is an antibody.

66. A pharmaceutical composition, comprising:

15

an expression vector encoding a polypeptide comprising the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:29; and

a pharmaceutically acceptable carrier.

20

67. The pharmaceutical composition of claim 66 wherein the expression vector comprises SEQ ID NO:1 or SEQ ID NO:28.

25

68. A method of treating a protein disulfide isomerase-like enzyme dysfunction related disease, wherein the disease is selected from cancer, diabetes and obesity, comprising the step of:

30

administering to a patient in need thereof a therapeutically effective dose of a reagent that modulates a function of a human protein disulfide isomerase-like enzyme, whereby symptoms of the protein disulfide isomerase-like enzyme dysfunction related disease are ameliorated.

- 84 -

69. The method of claim 68 wherein the reagent is identified by the method of claim 36.
- 5 70. The method of claim 68 wherein the reagent is identified by the method of claim 45.
71. The method of claim 68 wherein the reagent is identified by the method of claim 49.

Fig. 1

```
atgaagctca ccctagagtc          ttctctatag tccctgaaa aggcattctca ttggaagtgg
ttttattcaa aacttcagcg          agcattttaat cactgaagtg acaactgata ccttttggga
ctctgcccag ttcccgctctc          acgttctcct  gctctattac gctccgtggg  gcggttctg
agtagtcctt caaaaacagg          tcatccagct  agtcggaac ctgcccattgg acacattcac
tccatccctc aatcacatct          ctccgaatga  ccttccttgg gaatttatgg tcgatcgtct
tgtggcaagg attgacgtgt          cctgcaacag  aaaggacctc agtgtgaaat accccgaaga
tcctactgtc ttgttttttc          acctgttgag  gttcattttg catcactcag accctgcttc
cgccccccag aatgtggcta          actctcctac  caaggagtgt cttcagagcg aggcagtctt
acagcggggg cacatctccc          acttggagag  agagatccag aaactgagag cagaaataag
cagcctccag cgagcacaag          tgcaggtgga  gtcccagctc tccagtgcct ccacgcaca
gcaccggctg cggcagcagc          agcggggcct  ggaagagcag cacagcctgc agctggcccc
cagtgagcag ctgcaggccc          tctatgagca  gaagacacgt gagctgcagg accgagaaca cgtggctcaa
caagctgcag gagctggccc          atgcctcaga  agaggaaact ggagggcagg gatggagctg aaagcctggc
gatcctgggt gcgaccatgg          agagggaaact tgagccctca gccaccccc agctccctgg
ggcccagaga gaggtccacc          ccaagcagcc  atgtcagcgc tctgaaagga ataaggagaa
cagctcccct ccacctgcca          cagcacagac  taa
```

- 2/25 -

Fig. 2

MKLTLESFIQ NFSVLYSPLK RHLIGGSAQ FPSQHLITEV TTDTFWEVVL QKQDVLLLYY
 APWCGFCPSL NHIFIQLARN LPMDTFTVAR IDVSQNDLPW EFMVDRLLPTV LFFPCNRKDL
 SVKYPEDVPI TLPNLLRFIL HHSDPASSPQ NVANSPKTEC LQSEAVLQRG HISHLEREIQ
 KLRAEISSLQ RAQVQVESQL SSARRDEHRL RQQORALEEQ HSLLLHAHSEQ LQALYEQKTR
 ELQELARKLQ ELADASENLL TENTWLKILV ATMERKLEGR DGAESLAAQR EVHPKQPEPS
 ATPQLPGSSP PPANVSATLV SERKENRTD

Fig. 3

MKISNLLAAF LAFSGGFFCA SAEVPKVNKE GLNELITADK VLMVKFYAPW CGHCKALAPE
 YESAADELEK DGISLVEVDC TEEGDLCEY SIRGYPTLNV FKNGKQISQY SGPRKHDALV
 KYMRKQLLPT VKPISKDTLE NFVEKADDLA VVAFFKDQKL NDTYTEVAEV MKDDFVFAAS
 DDKELAKSLG SNFPGIVAFK KDAAQDSDKL VYTGDWDPAS IADFIGVSSI PLLDELNQMT
 FGKYQQSGLP LGIIFYNSTE SRDELYDVFFQ PLAKKYQDTL RFAFLDAVRY GAVAKQMNVE
 SDWPAFVIAN LKSMWKYPFP TTELTAKAMT KFGVDFVDGK LQPKIKSQPI PESQEDLVVL
 VADNFDDIVM DETKDVLEF YAPWCGHCKN LAPTYEKLAE EYSDDSNVVV AKIDATENDI
 SVSISGFPTI MFFKANDKVN PVRYEGDRTL EDLSAFIDKH ASFEPIKKEK ESWPAPDLED
 QVAVEDEMAD EL

Fig. 4

ssvvvlt denFdeevlkaksdkpVLv dFyApwCGpC

Fig. 5

```

ttttaacttc ctatagcccc ttcagctact acactgcatg ttgcaggacc ataagcaggg
gtgtgtcagg cttcatcgac tctgaacaag gtgtctttga agcccctact gttgcatttt
cttcccttga gaagaaatgt gaggttgatg cccaagctc cgttcctcac attgaggaga
acaggatatc ctttccagaa gtggacatga gtggacatga ctacacaaa cttcacaggc ctgagctgca
gaaccaacaa gactctcaac atctaccttt cactcaggt gaaagaattt gcggcaattg ttgacgtgaa
gagactgggt gctccgagct cactcaggt tggatccaaa gcaagcactg atgaagctca cctagagtc
agaagaatct cattacatct tggatccaaa tctctatag tccctgaaa aggcattctca ttggaagtgg
ttttattcaa aacttcagcg tctctatag agcatttaat cactgaagtg acaactgata ccttttggga
ctctgcccag ttcccgcttc acgttctcct gctctattac gctccgtggt gctccatgg acacattcac
agtagtcctt caaaaacagg tcatccagct ccttccttgg gaatttatgg accccgaaga
tccatccctc aatcacatct ctcagaatga aaaggacctc agtgtgaaat accctgcttc
tgtggcaagg attgacgtgt cctgcaacag gttcattttg catcactcag aggcagtctt
tcctactgtc ttgttttttc acctgttgag caaggagtgt aaactgagag cagaaataag
cgtccctatc acccttccaa actctctac agagatccag tccagtgcc cccacgcaca
cagccccccag aatgtggcta actctctac gttccagctc gcaagagcag cacagcctgc tccacgcaca
acagcggggg cacatctccc acttgagag agagatccag ggtccagctg gatggcctg
cagcctccag cgagcacaag tgcaggtgga gtccagctc ggaagagcag gaagacagt accgagctca
gcaccggctg cggcagcagc agcgggccc tctatgagca gaaacctcct accgagctca cgtggctcaa
cagtgagcag ctgcaggccc tctatgagca gaaacctcct accgagctca gatggcctg aaagcctggc
caagctgcag gagctggccg atgcctcaga aacacctcct accgagctca gatggcctg aaagcctggc
gatcctgggt gcgaccatgg agaggaact ggagggcagg tgagccctca gccaccccc agtccctgg
ggccagaga gaggtccacc ccaagcagcc tgagccctca tgagccctca gccaccccc agtccctgg
cagctccctt ccactgcca atgtcagcgc cacactgggt cactgaaagg tctgaaagg ataaggagaa
caggacagac taacttttta aatgatatga agaaatcaga ggtgaaaatt gtacattggg
aataatatata tgcaaaattt attgaaattt attgaaata agatttttct cagtgggtcta
gaaaatcaaa aaaaaaaaaa aaaaaaaaaa aa

```


Fig. 6

ttttaacttc ctatagcccc ttcagctact aactgcatg ttgcaggacc ataagcagggg
gtgtgtcagg cttcatcgac tctgaacaag gtgtccttga agccctact gttgcatttt
cttcccttga gaagaaatgt gaggtgatg cccaagctc cgttcctcac attgaggaga
acaggtatct ctttccagaa gtggacatga ctgacacaaa ctgcacaggc ctgagctgca
gaaccaaaa gactctcaac atctacctt tggattcaaa tttgtttggt tatatgcaga
gagactgggt gctccgagct ccactcaggt gaaagaattt gcggaattg ttgacgtgaa
agaagaatct cattacatct tggatccaaa gcaagcactg atgaagctca ccctagagtc
ttttattcaa aacttcagcg ttctctatag tccttgaaa aggcattctca ttggaagtgg
ctctgcccag ttcccgtctc agcatttaat cactgaagtg acaactgata ccttttggga
agtagtcctt caaaaacagg acgttctcct gctctattac gctccgtggt gcggtctctg
tccatccctc aatcacatct tcatccagct agtcggaac ctgcccattg acacattcac
tgtggcaagg attgacgtgt ctcagaatga cttcccttg gaatttatgg tcgacgtct
tcctactgtc ttgttttttc cctgcaacag aaaggaccta agtgtgaaat accccgaaga
cgtccctatc acccttccaa acctgttgag gttcattttg catcactcag accctgcttc
cagcccccag aatgtggcta actctcctac caaggagtgt cttcagagcg aggcagtctt
acagcggggg cacatctccc acttggagag agagatccag aaactgagag cagaaataag
cagcctccag cgagcacaag tgcagggtgga gtcccagctc tccagtgcc cccacgcaca
gcaccggctg cggcagcagc agcggggcct ggaagagcag cacagcctgc tccacgcaca
cagtgagcag ctgcaggccc tctatgagca gaagacacgt gagctgcagg agctggcccg
caagctgcag gagctggccg atgcctcaga aaacctcct accgagaaca cgtggctcaa
gatcctggtg gcgaccatgg agagggaaact ggagggcagg gatggagctg aaagcctggc
ggcccagaga gaggtccacc ccaagcagcc tgagccctca gccaccccc agctccctgg

Fig. 6 (continued)

cagctcccct ccacctgcca atgtcagcgc cacactggtg tctgaaagga ataaggagaa
caggacagac taacttttta aatgatatga agaaatcaga ggtgaaaatt gtacattggg
aatatattta tgcaaatttt attgaaattt attgtaaata aagattttct cagtgggtcta
gaaatcaaa aaaaaaaaaa aaaaaaaaaa aa

Fig. 7

```

ttttaacttc ctatagcccc ttcagctact acactgcatg ttgcaggacc ataagcaggg
gtgtgtcagg cttcatcgac tctgaacaag gtgtctttga agcccctact gttgcatttt
cttccttga gaagaaatgt gaggttgatg cccaagctc cgttcctcac attgaggaga
acaggtatct ctttccagaa gtggacatga ctagcacaaa ttgtttggt tatatgcaga
gaaccaacaa gacttcaac atctacctt cactcaggt gaaagaattt gggcaattg ttgacgtgaa
gagactgggt gctccgagct cattacatct tggatccaaa gcaagcactg atgaagctca ccctagagtc
agaagaatct ctttattcaa aacttcagcg ttctctatag tcccttgaaa aggcattctca ttggaagtgg
ctctgcccag ttcccgtctc agcattttaat cactgaagtg acaactgata ccttttggga
agtagtcctt caaaaacagg acgttctcct gctctattac gctccgtggt gcggttctg
tccatccctc aatcacatct tcatccagct agctcggaac ctgcccattg acacattcac
tgtggcaagg attgacgtgt ctcagaatga ccttccttgg gaatttatgg tcgatcgtct
tcctactgtc ttgttttttc cctgcaacag aaaggacctc agtgtgaaat accccgaaga
cgtccctatc acccttccaa acctgttgag gttcattttg catcactcag accctgcttc
cagccccag aatgtggcta acttcctac caaggagtgt cttcagagcg aggcagtctt
acagcggggg cacatctccc acttgagag agagatccag aaactgagag cagaaataag
cagcctccag cgagcacaag tgcaggtgga gtcccagctc tccagtgcgc gcagagatga
gcaccggctg cggcagcagc agcgggccc tctatgagca ggaagagcag cacagcctgc tccacgcaca
cagtgagcag ctgcaggccc tctatgagca gaagacacgt gagctgcagg agctggcccg
caagctgcag gagctggccg atgcctcaga aaacctcctt accgagaaca cgtggctcaa
gatcctgggt gcgaccatgg agaggaaact ggagggcagg gatggagctg aaagcctggc
ggcccagaga gaggtccacc ccaagcagcc tgagccctca gccaccccc agtccctgg

```

Fig. 7 (continued)

cagctcccct ccacctgccca atgtcagcgc cacactgggtg tctgaaagga ataaggagaa
caggacagac taacttttta aatgatatga agaaatcaga ggtgaaaatt gtacattggg
aatatattta tgcaaatattt attgaaattt attgtaaata aagattttct cagtgggtcta
gaaatcaaa aaaaaaaaaa aaaaaaaaaa aa

Fig. 8

ACCACTGAGAAAATCTTTATTACAATAAAATTTCAATAAAATTTGCATAAATATATATCCC
AATGTACAAATTTTCACTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCTGTTC
TCCTTATTCCTTTCAGACACCAAGTGTGGCGCTGACATTGGCAGGTGGAGGAGCTGCCA
GGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGTGGACCTCTCTCTGGGCCGCC
AGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCACAGGATCTTG
AGCCAGTGTCTCGGTAAGGAGGTTTCTGAGGCATCGGCCAGCTCCTGCAGCTTGCGG
GCCAGCTCCTGCAGCTCACGTGTCTTCTGCTCATAGAGGCCCTGCAGCTGCTCACTGTGT
GCGTGAGCAGGCTGTGCTGCTCTTCCAGGGCCCGCTGCTGCTGCCGAGCCGGTGCTCA
TCTCTCGGGCACTGGAGAGCTGGGACTCCACCTGCACTTGTGCTCGCTGAGGCTGCTT
ATTTCTGCTCTCAGTTTCTGGATCTCTCTCTCCAAGTGGGAGATGTGCCCCCGCTGTAAG
ACTGCCCTCGCTCTGAAGACACTCCTTGGTAGGAGAGTTAGCCACATTCTGGGGGCTGGAA
AGCAGGGTCTGAGTGATGCAAAATGAACCTCAACAGGTTTGGAAAGGTGATGGGACGTC
TTCGGGGTATTTACACACTTAGGGTCCCTTTCTGTTCAGG

Fig. 9

TAGACCACTGAGAAAATCTTTATTATTAACAATAAAATTTCAATAAAATTTGCATAAAATATATT
CCCAATGTACAAATTTTACCTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCTCTG
TTCTCCTTATTCTTTTCAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGGAGGGAGCTG
CCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGTGGACCTCTCTCTGGGCC
GCCAGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGCTGCTGCCACCCAGGATC
TTGAGCCACGTGTTCTCGGTAAGGAGTTTCTGAGGCATCGGCCAGCTCCTGCGAGCTTG
CGGCCAGCTCCTGCAGCTCACGTGCTCTCTGCTCATAGAGGGCTGCAGCTGCTCACTG
TGTGCGTGGAGCAAGCTGTGCTCTTCCAGGGCCGCTGCTGCTGCCGAGCCGGTGC
TCATCTCTGCGGCACTGGAGAGCTGGGACTCCACCTGCACTTGTGCTCGCTGGAGGCTG
CTTATTCTGCTCTCAGTTTCTGGATCTCTCTCCAAGTGGAGATGTGCCCCCGCTGT
AAGACTGCCCTCGCTCTGAAGACACTCCTTGGTAGGAGAGTTAGCCACATTTCTNNGGGCTG
GAAGCAGGGTCTGAATGATGCAAAATGAACCCCTAACAGGTTTGGAAAGTGATGGGGAGG
TCTTCCGGGGTATTTTACACTTAGGTCCCTTCTGTGGACGGAAAAAACAAGCG

Fig. 10

TAGACCACTGAGAAAATCTTTATTATTAACAATAAAATTTCAATAAAATTTGCATAAAATATATT
CCCAATGTACAAATTTTACCTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCTCTG
TTCTCCTTATTCTTTTCAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGGAGGGAGCTG
CCAGGAGCTGGGGGTGGCTGAGGGCTCANGCTGCTTGGGTGGACCTCTCTCTGGGCC
GCCAGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCACCCAGGATC
TTGAGCCACGTGTTCTCGGTAAGGAGTTTCTGAGGCATCGGCCAGCTCCTGCAGCTTG
CGGGCCAGCTCCTGCAGCTCACGTGCTCTTCTGCTCATAGAGGCCCTGCAGCTGCTCACTG
TGTGCGTGGAGCAGGCTGTGCTGCTCTTCCAGGGCCGCTGCTGCTGCCGAGCCGGTGC
TCATCTCTGCGGGCACTGGAGAGCTGGGACTCCACCTGCACTTGTGCTCGCTG

- 9/25 -

Fig. 11

CGAGCTCCACTCAGGTGAAAGAAATTGCGGCAATTGTTGACGTGAAAGAAATCTCATT
ACATCTTGGATCCAAAGCAAGCACTGATGAAGCTCACCTAGAGTCTTTTATTCAAACCT
TCAGCGTTCTCTATAGTCCCTTGAAAGGCACTCTCATTTGGAAGTGGCTCTGCCAGTTCC
CGTCTCAGCATTTAATCACTGAAGTGACAACTGATACCTTTTGGGAAGTAGTCTTCAAA
AACAGGACGTTCTCCTGCTCTATTACGCTCCGTGGTGGCTTCTGTCCATCCCCTCAATC
ACATCTTCATCCAGCTAGCTCGGAACCTGCCCATGGACACATTCACTGTGGCAAGGATTG
ACGTGCTCTCANAATGACCTTCTTTGGGAATTATGGTCGATCGTCNCCFACTGCTTGT
TTTTTCCCTGCAACAGAAAGGACCTAAGTGTGAAATACCCGGAAGACGTCTCATCACCC
TTCCAAACCTGGTGAGGTTCAATTTTGCACTCACTCAGACCCCTGCTTCCAGCCCCCAGATGT
GGC

Fig. 12

TAGACCACTGAGAAAATCTTTATTACAATAAATTTCAATAAAAATTTGCATAAATATATT
CCCAATGTACAATTTTCACCTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCCCTG
TTCTCCTTATTCCTTTCAGACACCAAGTGTGGCGCTGACATTGGCAGGTGGAGGGAGCTG
CCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGGACCTCTCTCTGGGCC
GCCAGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCACAGGATC
TTGAGCCACGTGTTCTCGGTAAGGAGGTTTCTGAGGCATCGGCCAGCTCCTGCAGCTTG
CGGGCCAGCTCCTGCAGCTCACGTGCTTCTGCTCATAGAGGGCTGCAGCTGCTCACTG
TGTGCGTGGAGCAGGCTGTGCTGCTCTTCCAGGGCCCCGCTGCTGCTGCCGAGCCGGTGC
TCATCTCTGCG

- 10/25 -

Fig. 13

CTGAAGTGACAACTGATACCTTTTGGGAAGTAGTCCTTCAAAAACAGGACGTTCTCCTGC
TCTATTACGCTCCGTGGTGGGCTTCTGTCCATCCCTCAATCACATCTTCAATCCAGCTAG
CTCGGAACCTGCCCATGGACACATTCACGTGGCAAGGATTGACGTGTCTCAGAATGACC
TTCCCTTGGGAATTATGGTCGATCGTCTTCCTACTGTCTTGTCTTTTCCCTGCAACAGAA
AGGACCTAAGTGTGAAATACCCGGAAGACGTCCCCATCACCTTCCAAACCTGTTGAGGT
TCATTTTGCATCACTCAGACCTTGCTCCAGCCCCCAGAAATGTGGCTAACTCTCCTACCA
AGGAGTGTCTTCAGAGCGAGGCAGTCTTACAGCGGGGGCACATNTNCCACTTGGAGAGAG
AGATCCAGAACTGAGAGCAGAAATAAGCAGCTNCAGCGAGCACAAAGTGCAGGTGGAGT
CCCAGCTC

Fig. 14

CTAGACCACTGAGAAAATCTTTATTTACAATAAAATTTCAATAAAAATTTGCATAAATATAT
TCCCAATGTACAATTTTCACTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCTT
GTTCTCCTTATTCTTTTCAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGGAGGGAGCT
GCCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGGACCTCTCTCTGGGC
CGCCAGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCACCCAGGAT
CTTGAGCCACGTGTTCTCGGTAAGGAGGTTTCTGAGGCATCGGCCAGCTCCTGCAGCTT
GCGGCCAGCTCCTGCAGCTCACGTGTCTTCTGCTCATAGAGGGCCTGCAGCTGCTCACT
GTGTGCGTGGAGCAGGCTGTGCTGCTCTTCCAGGGCCCGCTGCTGCTG

Fig. 15

TTTTTTNATGTTTTTTTAGACCACTGAGAAAATCTTTATTACAAATAAAATTTCAATAAA
ATTTGCATAAATATATTTCCCAATGTACAATTTTACCTCTGATTTCTTCATATCATTTAA
AAAGTAGTCTGTCCTGTTCTCTTATTCTTTTACAGACACCAGTGTGGCGCTGACATTGG
CAGGTGAGGGAGCTGCCAGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGT
GGACCTCTCTCTGGCCCGCAGGCTTTTACGCTCCATCCCTGCCCTCCAGTTTCCCTCTCCA
TGGTCGCCACCAAGATCTTGAGCCACGTGTTCTCGGTAGGAGGTTTTCTGAGGCATCGG
CCAGTCTCTGCAGCTTGCGGGCCAGCTCCTGCAGCTCACGTGCTTCTTGCTCATAGAGGG
CCTGCAGCTGCTCACTGTGTGCGTGAGCAGGCTGTGCT

Fig. 16

TCTAGACCACTGAGAAAATCTTTATTACAAATAAAATTTCAATAAAATTTGCATAAATA
TTCCCAATGTACAATTTTACCTCTGATTTCTTCATATCATTTAAAGTTAGTCTGTCC
TGTTCTCCTTATTCCTTTCAGACACCAGTGTGGCGCTGACATTGGCAGGTGGAGGGAGC
TGCCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGACCTCTCTCTGGG
CCGCCACGGTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCACCAGGA
TCTTGAGCCACGTGTTCTCCGTAGAGNNTTTCTGAGGCATCGGCCAGCTCCTGCAGCT
TGCGAGCCAGCTCCTGCAGCTCACGTGCTTCTTGCTCATAGAGGGCTGCAGCTGCTCAC
TGTGTGCGTGAGCAGGCTGTGTGCTGCTCTTCCAGGGCCCGCTGCTGCTGCCG

Fig. 17

TCTAGACCACTGAGAAAATCTTTATTACAAATAAATTTCAATAAAATTTGCATAAATATA
TTCCCAATGTACAAATTTTCACTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCC
TGTTCTCCTTATCCTTTAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGGAGGGGAGC
TGCCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGTGGACCTCTCTCTGGG
CCGCCAGGCTTTGAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCCGCCACCAGGA
TCTTGAGCCACGTGTCTCGGTAAGGAGGTTTCTGAGGCATCGGCCAGCTCCTGCAGCT
TGCGGGCCAGCTCCTGCAGCTCACGTGTCTTCTGTCTCATAGAGGGCCTGCAGCTGCTCAC
TGTGTGCGTGGAGC

Fig. 18

GATTTTCTAGACCACTGAGAAAATCTTTATTACAAATAAATTTCAATAAAATTTGCATAA
ATATATTCCCAATGTACAAATTTTCACTCTGATTTCTTCATATCATTTAAAAAGTTAGTC
TGTCTGTCTCCTTATCCTTTAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGGAGG
GGAGCTGCCAGGGAGCTGGGGGTGGCTGAGGGCTCANGCTGCTTNGGTGGACCTNTCT
CTGGGCCGCCAGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCCGCCAC
CAGGATCTTGAGCCACGTGTCTCGGTAAGGAGGTTTCTGAGGCATCGGCCAGCTCCTG
CAGCTTGCGGGCCAGCTCCTGCAGCTCACGTGTCTTCTGTCTCATAGAGGGCCTGCAGCTG
CTCACTGTGTGCGTGGAGCA

Fig. 19

AGACCATTTAGTAAATCTTTATTACAAATAAATTTCAATGAAATTTGCATAAAATATATTC
CCAATGTACAAATTTTCACTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCTGT
TCTCCTTATTCCTTTCAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGGAGGGAGCTGC
CAGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGTGGACCTCTCTCTGGGCCG
CCAGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGTCGCCACCCAGGATCT
TGAGCCACGTGTTCTCGGTAAGGAGGTTTCTGAGGCATCGGCCAGTCTCTGCAGCTTGC
GGCCAGCTCCTGCAGCTCACGTGCTCTCTGCTCATAGAGGGCCTGCAGCTGCTCACTGT
GTGCGTGAGCAGGCTGTGCTGCTCT

Fig. 20

TTTTTTTTTCTAGACCACTGAGAAATCTTTATTTACAAATAAATTTCAATAAAATTTGCA
TAAATATATTTCCCAATGTACAAATTTTCACTCTGATTTCTTCATATCATTTAAAAAGTTA
GTCTGTCTGTTCTCCTTATTCCTTTCAGACACCAAGTGTGGCGCTGACATTTGCAGGTGG
AGGGAGCTGCCAGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGGACCTC
TCTCTGGCGCCACGTTTCAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCC
ACCAGGATCTTGAGCCACGTGTTCTCGGTAAGGAGGTTTCTGAGGCATCGGCCAGCTCC
TGCAGCTTGCGGGCCAGCTCCTGCAGCTCACGTGCTCTTCTGCTCATAGAGGGCCTGCAGC
TGCTCACTGTGTGCGTGGAG

Fig. 21

TTTTTTTTCTAGACCACCTGAGAAAATCTTTATTACAAATAAAATTTCAATAAAATTTGCA
TAAATATATCCCAATGTACAATTTTCAACCTCTGATTTCTTCATATCATTTAAAAAGTTA
GTCTGCTCTGTTCTCCTTATTCTTTTACAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGG
AGGGAGCTGCCAGGAGCTGGGGGTGGCTGAGGCTCAGGCTGCTTGGGGTGGACCTC
TCTCTGGGCGCCACGTTTACAGCTCCATCCCTGCCCTCCAGTTTCCCTCTCCATGGTCCG
CACCAGGATCTTGAGCCACGTGTTCTCGGTAAGGAGGTTTCTGAGGCCATCGGCCAGCTC
CTGCAGCTTGCGGGCCAGCTCCTGCAGCTCACGTGTCTTCTGCTCATAGAGGGCCTGCAG
CTGCTCACTGTGTGCGTGGAG

Fig. 22

CTAGACCACCTGAGAAAATCTTTATTACAAATAAAATTTCAATAAAATTTGCATAAAATAT
TCCCAATGTACAATTTTCAACCTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCTT
GTTCTCCTTATTCTTTCAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGGAGGAGCT
GCCAGGAGCTGGGGGTGGCTGAGGCTCAGGCTGCTTGGGGTGGACCTCTCTCTGGGC
CGCCACGTTTACAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCACCAGGATC
TTGAGCCACGTGTTCTCGGTAAGGAGGTTTCTGAGGCCATCGGCCAGCTCCTGCAGCTTG
CGTGCCAGCTCCTGCAGCTCACGTGTCTTCTGCTCATAGAGGGCCTGCAGCTGCTCACTG
TGTGCGTGGAGCAGGCT

Fig. 23

TTTTCTAGACCACTGAGAAAATCTTTATTACAATAAAATTTCAATAAAAATTTGCATAAAT
ATATTCCCAATGTACAATTTTCACCTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTG
TCCTGTTCTCCTTATTCCCTTTCAGACACCAAGTGTGGCGCTGACATTGGCAGGTGGAGGGG
AGTGCCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGGACCTCTCTCT
GGCCGCCACGTTATCAGCCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCACC
AGGATCTTGAGCCACGTGTTCTCGGTAAGGAGTTTCTGAGGCATCGGCCAGCTCCTGC
AGCTTGCGAGCCAGCTCCTGCAGCTCACGTGCTTCTGCTCATAGAGGGCCTGCAGCTGC
TCACTGTGTGCGTGGAG

Fig. 24

TTTTTTCTAGACCACTGAGAAAATCTTTATTACAATAAAATTTCAATAAAAATTTGCATAA
ATATATTCCCAATGTACAATTTTCACCTCTGATTTCTTCATATCATTTAAAAAGTTAGTC
TGTCTGTTCTCCTTATTCCCTTTCAGACACCAAGTGTGGCGCTGACATTGGCAGGTGGACG
GGAGCAGCCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGGACCTCTCT
CTGGCCGCCACGTTCTAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGCCAC
CAGGATCTTGAGCCACGTGTTCTCGGTAAGGAGTTTCTGAGGCATCGGCCAGCTCCTG
CAGCTTGCGGGCCAGCTCCTGCAGCTCACGTGCTTCTGCTCATAGAGGGCCTGAAGCTG
CTCACTGTGTGCGTGGAGAAG

Fig. 25

TTTTTTTTTTCTAGACCACTGAGAAAATCTTTATTACAAATAAAATTTCAATAAAA
TTTGCATAAATAATATCCCAATGTACAATTTTACCTCTGATTTCTTCATATCATTTAAA
AAGTAGTCTGTCTCTCTTATTCCTTTTACAGACACCAAGTGTGGCGCTGACATTTGGC
AGGTGAGGGAGCTGCCAGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTG
GACCTCTCTCTGGGCCGCCAGGCTTTTACAGCTCCATCCCTGCCCTCCAGTTTCTCTCCAT
GGTCGCCACCAGGATCTTGAGCCACGTGTTCTCGGTAAGGAGGTTTCTGAGGCATCGGC
CAGCTCCTGCAGCTTGCGGGCCAGCTCCTGCAGCTCACGTGTCTTCTGCTCATA

Fig. 26

TTTTTTTTTTCTAGACCACTGAGAAAATCTTTATTACAAATAAAATTTCAATAAAATTTGCA
TAAATATATCCCAATGTACAATTTTACCTCTGATTTCTTCATATCATTTAAAAGTTA
GTCTGTCTGTCTCTCTTATTCCTTTTACAGACACCAAGTGTGGCGCTGACATTTGGCAGGTGG
AGGGAGCTGCCAGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGGACCTC
TCTCTGGGCCGCCAGGCTTTTACAGCTCCATCCCTGCCCTCCAGTTTCTCTCCATGGTCGC
CACCANGATCTTGAGCCACGTGTTCTCGGTAGGAGGTTTCTGAGGCATCGGCCAGCTC
CTGCAGCTTGCGGGCCAGCTCCTGCACTTCAACGTGCTCTCTGCTCATTAGAGGGCCTGC
AGCTGCTCACTGTGTGCGGTGGAGCAAGGCCTGTGCTTG

Fig. 27

CTAGACCACTGAGAAAATCTTTATTACAAATAAAATTTCAATAAAATTTGCATAAAATATAT
TCCCAATGTACAATTTTACACCTCTGATTTCTTCATATCATTTAAAAAGTTAGTCTGTCTCT
GTTCTCCTTATTCCCTTTCAGACACACAGTGTGGCGCTGACATTGGCAGGTGGAGGGAGCT
GCCAGGGAGCTGGGGGTGGCTGAGGGCTCAGGCTGCTTGGGGTGGACCTCTCTCTGGGC
CGCAGGGCTTTCAGCTCCATCCCTGCCCTCCAGTTTCCCTCTCCATGGTCGCCACCCAGGAT
CTTGAGCCACGTGTTCTCGGTAAGGAGGTTTCTTGAGGCATCGGCCAGCTCCTTGCAGCTT
GCGGGCCAGCTCCTGCAGCTCACGTGTCTTCTGTCTCATA

- 18/25 -

FIG. 28

BLASTP - alignment of 59_PROTEIN (SEQ ID NO:2) against swiss|Q10057|PDI1_SCHPO (SEQ ID NO:3)

This hit is scoring at : 7e-10 (expectation value), Alignment length (overlap): 131, Identities : 32 %, Scoring matrix : BLOSUM62 (used to infer consensus pattern), Database searched : nrdb

Q: 33 SQHLITEVTTDTFWEVVLQK-QDVLLLYAPW**CGFC**PSLNHIFIQLARNLPMDT-FTVAR
 SQ: .. :..D.F ::V::: :DVL: :YAPWCG.C.:L :: :LA.... D: ..VA:
 H: 353 SQEDLVVLVADNFDDIVMDETKDVLVEFYAPW**CGHC**CKNLAPT^YEKLAEYSDDSNVVVAK

disulfe redox-active site CXXC motif

BLOCKS for Thio redoxin family proteins

IDVSQNDLPWEFMVDRRLPTVLFFPCNRKDL^SVKYPEDVPITLPNLLRFILHHS--DPASS
 ID::ND:PT::FF..N K .V:Y. D TL.:L .FI .H: :P...
 IDATENDI--SVSISGFPTIMFFKANDKVN^PVR^YEGD--RTLEDLSAFIDKHASFEPIKK

PQNVANSPTKE 159

... :P. E

EKESVPAPDLE 479

- 19/25 -

FIG. 29

BLOCKS search results

AC#	Description	Strength	Score
BL00194	Thioredoxin family proteins.	1242	1223
AA# 55	LLLYAPWCGfCP		

- 20/25 -

FIG. 30

HMMPFAM - alignment of 59_PROTEIN (SEQ ID NO:2) against pfam|hmm|thioredoxin (SEQ ID NO:4)

Thioredoxin

This hit is scoring at : 26.4; Expect = 1.2e-06, Scoring matrix : BLOSUM62 (used to infer consensus pattern),

Q:	34 QHLITEVTDTFWEVVLQ---KQDVLLLLYYAPWCGFC	67
H:	1 ssvvvltDenFdeevlkaksdKpVLVdFyApWCGpC	37

Fig. 31

atgaagctcaccctagagtcttttattcaaaacttcagcgttctctatagtcaccttgaaa
aggcatctcattggaagtggctctgcccagttcccgtctcagcatttaatactgaagtg
acaactgataaccttttgggaagtagtccctcaaaaacaggacgttctcctgctctattac
gctccgtggcggtctctgtccatccctcaatcacatcttcacccagctagctcggaac
ctgcccattggacacattcactgtggcaaggattgacgtgtctcagaatgaccttccttgg
gaatttatggtcgatcgtcttcctactgtctgtttttccctgcaacag

Fig. 32

MKLTLESFIQ NFSVLYSPLK RHLIGSGSAQ FPSQHLITEV TTDTFWEVVL QKQDVLLLYY
APWCGFCPSL NHIFIQLARN LPMDTFTVAR IDVSQNDLPW EFMVDRLPTV LFFPCN

Fig. 33

BLASTP - alignment of 59_short_TR1 against swiss|P30101|ER60_HUMAN
This hit is scoring at : 4e-09 (expectation value), Alignment length (overlap)
: 116, Identities : 33 %, Scoring matrix : BLOSUM62 (used to infer consensus
pattern), Database searched : nrdb

```

Q:      5  LESFIQNFVLYSPLKRHLIGSGSAQFPSQH--LITEVTDTTFWEVVL-QKQDVLLLYYA
      LE.F:Q::  . LKR:L  S...P...  :. V....F E:V  :.:DVL: :YA
H:     349  LERFLQDY--FDGNLKRYL---KSEPIPESNDGPVKVVAENFDEIVNNENKDVLIIFYA

      PWC GFCPSLNHIFIQLARNLPMD-TFTVARIDVSQNDLPWEFMVDR LPTVLFFPCN      116
      PWC G.C.:L.  : :L...L..D ...:A::D::ND:P .: V  .PT::F P.N
      PWC GCHCKNLEPKYKELGKLSKDPNIVIAKMDATANDVPSPYEVVRGFPFTIYFSPAN      459
disulfe redox-active site CXXC motif
  
```

BLOCKS for Thioredoxin family proteins

AC#	Description	Strength	Score
BL00194	Thioredoxin family proteins.	1242	1223
AA# 55	LLLYYAPWCGfCP		

Fig. 34


HMMPFAM - alignment of 59_short_TR1 against pfam|hmm|thioered
Thioredoxin, This hit is scoring at : 26.4; Expect = 1.2e-06, Scoring matrix :
BLOSUM62 (used to infer consensus pattern)

Q:	34	QHLITEVTTFWEVVLQ---	KQDVLLLLYYAPWCGFC	67
H:	1	ssvvvlt denFdeevlkaksd	kpVLVdFyApWCGpC	37

Fig. 35

Sample	Cell Name	Abs. #	Norm. #
Sample 1	Brain-1	1579	1579
Sample 2	Heart-3	1774	3042
Sample 3	Kidney	6828	8885
Sample 4	Liver	3558	11115
Sample 5	Lung	8503	17180
Sample 6	Trachea	4762	7600
Sample 7	Bone Marrow-1	1161	1976
Sample 8	Colon	4338	3031
Sample 9	Small intestine	3784	5051
Sample 10	Spleen-1	4244	5601
Sample 11	Stomach	4511	5420
Sample 12	Thymus-1	2646	2579
Sample 13	Mammary gland	4032	6035
Sample 14	Prostate-1	2645	2271
Sample 15	Skeletal muscle-1	7737	12329
Sample 16	Testis	9153	7092
Sample 17	Uterus	2286	5038
Sample 18	Cerebellum	5786	5111
Sample 19	Fetal Brain	4418	2603
Sample 20	Fetal Liver-1	1910	5728
Sample 21	Spinal cord	1792	1815
Sample 22	Placenta-1	5558	6713
Sample 23	Adrenal gland	2546	2480
Sample 24	Pancreas-1	1905	11854
Sample 25	Salivary gland	5268	7860
Sample 26	Thyroid	6422	11625

Fig. 36



SEQUENCE LISTING

<110> Bayer AG

Bull, Christof

<120> REGULATION OF HUMAN PROTEIN DISULFIDE ISOMERASE-LIKE ENZYME

<130> Lio132 foreign countries

<150> 60/231,297

<151> 2000-09-08

<160> 29

<170> PatentIn version 3.1

<210> 1

<211> 993

<212> DNA

<213> Homo sapiens

<400> 1

atgaagctca	ccctagagtc	ttttattcaa	aacttcagcg	ttctctatag	tcccttgaaa	60
aggcatctca	ttggaagtgg	ctctgcccag	ttcccgctctc	agcatttaat	cactgaagtg	120
acaactgata	ccttttgagg	agtagtcctt	caaaaacagg	acgttctcct	gctctattac	180
gctccgtggg	gcggtttctg	tccatccctc	aatcacatct	tcatccagct	agctcggaac	240
ctgcccattg	acacattcac	tgtggcaagg	attgacgtgt	ctcagaatga	ccttccttgg	300
gaatttatgg	tcgatcgtct	tcctactgtc	ttgttttttc	cctgcaacag	aaaggacctc	360
agtgtgaaat	accccgaaga	cgtccctatc	acccttccaa	acctgttgag	gttcattttg	420
catcactcag	accctgcttc	cagcccccag	aatgtggcta	actctcctac	caaggagtgt	480
cttcagagcg	aggcagtctt	acagcggggg	cacatctccc	acttgagag	agagatccag	540
aaactgagag	cagaaataag	cagcctccag	cgagcacaag	tgaggtgga	gtcccagctc	600
tccagtggcc	gcagagatga	gcaccggctg	cggcagcagc	agcgggccct	ggaagagcag	660
cacagcctgc	tccacgcaca	cagtgcagc	ctgcaggccc	tctatgagca	gaagacacgt	720
gagctgcagg	agctggcccc	caagctgcag	gagctggccg	atgcctcaga	aaacctcctt	780
accgagaaca	cgtggctcaa	gatcctgggt	gcgaccatgg	agaggaaact	ggagggcagg	840
gatggagctg	aaagcctggc	ggcccagaga	gaggtccacc	ccaagcagcc	tgagccctca	900

gccaccccc agctccctgg cagctcccct ccacctgcca atgtcagcgc cacactgggtg 960
 tctgaaagga ataaggagaa caggacagac taa 993

<210> 2

<211> 330

<212> PRT

<213> Homo sapiens

<400> 2

Met Lys Leu Thr Leu Glu Ser Phe Ile Gln Asn Phe Ser Val Leu Tyr
 1 5 10 15
 Ser Pro Leu Lys Arg His Leu Ile Gly Ser Gly Ser Ala Gln Phe Pro
 20 25 30
 Ser Gln His Leu Ile Thr Glu Val Thr Thr Asp Thr Phe Trp Glu Val
 35 40 45
 Val Leu Gln Lys Gln Asp Val Leu Leu Leu Tyr Tyr Ala Pro Trp Cys
 50 55 60
 Gly Phe Cys Pro Ser Leu Asn His Ile Phe Ile Gln Leu Ala Arg Asn
 65 70 75 80
 Leu Pro Met Asp Thr Phe Thr Val Ala Arg Ile Asp Val Ser Gln Asn
 85 90 95
 Asp Leu Pro Trp Glu Phe Met Val Asp Arg Leu Pro Thr Val Leu Phe
 100 105 110
 Phe Pro Cys Asn Arg Lys Asp Leu Ser Val Lys Tyr Pro Glu Asp Val
 115 120 125
 Pro Ile Thr Leu Pro Asn Leu Leu Arg Phe Ile Leu His His Ser Asp
 130 135 140
 Pro Ala Ser Ser Pro Gln Asn Val Ala Asn Ser Pro Thr Lys Glu Cys
 145 150 155 160
 Leu Gln Ser Glu Ala Val Leu Gln Arg Gly His Ile Ser His Leu Glu
 165 170 175
 Arg Glu Ile Gln Lys Leu Arg Ala Glu Ile Ser Ser Leu Gln Arg Ala
 180 185 190
 Gln Val Gln Val Glu Ser Gln Leu Ser Ser Ala Arg Arg Asp Glu His
 195 200 205
 Arg Leu Arg Gln Gln Gln Arg Ala Leu Glu Glu Gln His Ser Leu Leu
 210 215 220
 His Ala His Ser Glu Gln Leu Gln Ala Leu Tyr Glu Gln Lys Thr Arg
 225 230 235 240
 Glu Leu Gln Glu Leu Ala Arg Lys Leu Gln Glu Leu Ala Asp Ala Ser
 245 250 255
 Glu Asn Leu Leu Thr Glu Asn Thr Trp Leu Lys Ile Leu Val Ala Thr
 260 265 270
 Met Glu Arg Lys Leu Glu Gly Arg Asp Gly Ala Glu Ser Leu Ala Ala
 275 280 285
 Gln Arg Glu Val His Pro Lys Gln Pro Glu Pro Ser Ala Thr Pro Gln
 290 295 300

Leu Pro Gly Ser Ser Pro Pro Pro Ala Asn Val Ser Ala Thr Leu Val
 305 310 315 320
 Ser Glu Arg Asn Lys Glu Asn Arg Thr Asp
 325 330

<210> 3

<211> 492

<212> PRT

<213> Homo sapiens

<400> 3

Met Lys Ile Ser Asn Leu Leu Ala Ala Phe Leu Ala Phe Ser Gly Gly
 1 5 10 15
 Phe Phe Cys Ala Ser Ala Glu Val Pro Lys Val Asn Lys Glu Gly Leu
 20 25 30
 Asn Glu Leu Ile Thr Ala Asp Lys Val Leu Met Val Lys Phe Tyr Ala
 35 40 45
 Pro Trp Cys Gly His Cys Lys Ala Leu Ala Pro Glu Tyr Glu Ser Ala
 50 55 60
 Ala Asp Glu Leu Glu Lys Asp Gly Ile Ser Leu Val Glu Val Asp Cys
 65 70 75 80
 Thr Glu Glu Gly Asp Leu Cys Ser Glu Tyr Ser Ile Arg Gly Tyr Pro
 85 90 95
 Thr Leu Asn Val Phe Lys Asn Gly Lys Gln Ile Ser Gln Tyr Ser Gly
 100 105 110
 Pro Arg Lys His Asp Ala Leu Val Lys Tyr Met Arg Lys Gln Leu Leu
 115 120 125
 Pro Thr Val Lys Pro Ile Ser Lys Asp Thr Leu Glu Asn Phe Val Glu
 130 135 140
 Lys Ala Asp Asp Leu Ala Val Val Ala Phe Phe Lys Asp Gln Lys Leu
 145 150 155 160
 Asn Asp Thr Tyr Thr Glu Val Ala Glu Val Met Lys Asp Asp Phe Val
 165 170 175
 Phe Ala Ala Ser Asp Asp Lys Glu Leu Ala Lys Ser Leu Gly Ser Asn
 180 185 190
 Phe Pro Gly Ile Val Ala Phe Thr Lys Asp Ala Ala Gln Asp Ser Asp
 195 200 205
 Lys Leu Val Tyr Thr Gly Asp Trp Asp Pro Ala Ser Ile Ala Asp Phe
 210 215 220
 Ile Gly Val Ser Ser Ile Pro Leu Leu Asp Glu Leu Asn Gln Met Thr
 225 230 235 240
 Phe Gly Lys Tyr Gln Gln Ser Gly Leu Pro Leu Gly Ile Ile Phe Tyr
 245 250 255
 Asn Ser Thr Glu Ser Arg Asp Glu Leu Tyr Asp Val Phe Gln Pro Leu
 260 265 270
 Ala Lys Lys Tyr Gln Asp Thr Leu Arg Phe Ala Phe Leu Asp Ala Val
 275 280 285

Arg Tyr Gly Ala Val Ala Lys Gln Met Asn Val Glu Ser Asp Trp Pro
 290 295 300
 Ala Phe Val Ile Ala Asn Leu Lys Ser Met Leu Lys Tyr Pro Phe Pro
 305 310 315 320
 Thr Thr Glu Leu Thr Ala Lys Ala Met Thr Lys Phe Val Gly Asp Phe
 325 330 335
 Val Asp Gly Lys Leu Gln Pro Lys Ile Lys Ser Gln Pro Ile Pro Glu
 340 345 350
 Ser Gln Glu Asp Leu Val Val Leu Val Ala Asp Asn Phe Asp Asp Ile
 355 360 365
 Val Met Asp Glu Thr Lys Asp Val Leu Val Glu Phe Tyr Ala Pro Trp
 370 375 380
 Cys Gly His Cys Lys Asn Leu Ala Pro Thr Tyr Glu Lys Leu Ala Glu
 385 390 395 400
 Glu Tyr Ser Asp Asp Ser Asn Val Val Val Ala Lys Ile Asp Ala Thr
 405 410 415
 Glu Asn Asp Ile Ser Val Ser Ile Ser Gly Phe Pro Thr Ile Met Phe
 420 425 430
 Phe Lys Ala Asn Asp Lys Val Asn Pro Val Arg Tyr Glu Gly Asp Arg
 435 440 445
 Thr Leu Glu Asp Leu Ser Ala Phe Ile Asp Lys His Ala Ser Phe Glu
 450 455 460
 Pro Ile Lys Lys Glu Lys Glu Ser Val Pro Ala Pro Asp Leu Glu Asp
 465 470 475 480
 Gln Val Ala Val Glu Asp Glu Met Ala Asp Glu Leu
 485 490

<210> 4

<211> 37

<212> PRT

<213> Homo sapiens

<400> 4

Ser Ser Val Val Val Val Leu Thr Asp Glu Asn Phe Asp Glu Glu Val
 1 5 10 15
 Leu Lys Ala Lys Ser Asp Lys Pro Val Leu Val Asp Phe Tyr Ala Pro
 20 25 30
 Trp Cys Gly Pro Cys
 35

<210> 5

<211> 1532

<212> DNA

<213> Homo sapiens

<400> 5

ttttaacttc	ctatagcccc	ttcagctact	acactgcatg	ttgcaggacc	ataagcaggg	60
gtgtgtcagg	cttcacgcac	tctgaacaag	gtgtccttga	agcccctact	gttgcatctt	120
cttcccttga	gaagaaatgt	gagggttgatg	ccccaaagctc	cgttcctcac	attgaggaga	180
acaggtatct	ctttccagaa	gtggacatga	ctagcacaaa	cttcacaggc	ctgagctgca	240
gaaccaacaa	gactctcaac	atctaccttt	tggattcaaa	tttgtttggt	tatatgcaga	300
gagactgggt	gctccgagct	ccactcaggt	gaaagaattt	gcggcaattg	ttgacgtgaa	360
agaagaatct	cattacatct	tggatccaaa	gcaagcactg	atgaagctca	ccctagagtc	420
ttttattcaa	aacttcagcg	ttctctatag	tcccttgaaa	aggcatctca	ttggaagtgg	480
ctctgcccag	ttcccgtctc	agcattttaat	cactgaagtg	acaactgata	ccttttgggg	540
agtagtcctt	caaaaacagg	acgtttctcct	gctctattac	gctccgtggg	gcggcttctg	600
tccatccctc	aatcacatct	tcattccagct	agctcgggac	ctgcccattg	acacattcac	660
tgtggcaagg	attgacgtgt	ctcagaatga	ccttccttgg	gaatttatgg	tcgacgtctc	720
tcctactgtc	ttgttttttc	cctgcaacag	aaaggaccta	agtgtgaaat	accccgaaag	780
cgctccctatc	acccttccaa	acctgttgag	gttcattttg	catcactcag	accctgcttc	840
cagccccccag	aatgtggcta	actctcctac	caaggagtgt	cttcagagcg	aggcagtctt	900
acagcggggg	cacatctccc	acttgagag	agagatccag	aaactgagag	cagaaataag	960
cagcctccag	cgagcacaag	tgcaggtgga	gtcccagctc	tccagtggcc	gcagagatga	1020
gcaccggctg	cggcagcagc	agcggggcct	ggaagagcag	cacagcctgc	tccacgcaca	1080
cagtgcagcag	ctgcaggccc	tctatgagca	gaagacacgt	gagctgcagg	agctggcccg	1140
caagctgcag	gagctggccg	atgcttcaga	aaacctcctt	accgagaaca	cgtggctcaa	1200
gatectgggtg	gcgaccatgg	agaggaaact	ggagggcagg	gatggagctg	aaagcctggc	1260
ggcccagaga	gaggtccacc	ccaagcagcc	tgagccctca	gccaccccc	agctccctgg	1320
cagctccctc	ccacctgcca	atgtcagcgc	cacactgggtg	tctgaaagga	ataaggagaa	1380
caggacagac	taacttttta	aatgatatga	agaaatcaga	ggtgaaaatt	gtacattggg	1440
aatatattta	tgcaaatttt	attgaaattt	attgtaaata	aagattttct	cagtgggtcta	1500
gaaaatcaaa	aaaaaaaaaa	aaaaaaaaaa	aa			1532

<210> 6

<211> 1532

<212> DNA

<213> Homo sapiens

<400> 6

ttttaacttc	ctatagcccc	ttcagctact	acactgcatg	ttgcaggacc	ataagcaggg	60
gtgtgtcagg	cttcacgcac	tctgaacaag	gtgtccttga	agcccctact	gttgcatctt	120
cttcccttga	gaagaaatgt	gagggttgatg	ccccaaagctc	cgttcctcac	attgaggaga	180
acaggtatct	ctttccagaa	gtggacatga	ctagcacaaa	cttcacaggc	ctgagctgca	240
gaaccaacaa	gactctcaac	atctaccttt	tggattcaaa	tttgtttggt	tatatgcaga	300
gagactgggt	gctccgagct	ccactcaggt	gaaagaattt	gcggcaattg	ttgacgtgaa	360
agaagaatct	cattacatct	tggatccaaa	gcaagcactg	atgaagctca	ccctagagtc	420
ttttattcaa	aacttcagcg	ttctctatag	tcccttgaaa	aggcatctca	ttggaagtgg	480
ctctgcccag	ttcccgtctc	agcattttaat	cactgaagtg	acaactgata	ccttttgggg	540
agtagtcctt	caaaaacagg	acgtttctcct	gctctattac	gctccgtggg	gcggcttctg	600
tccatccctc	aatcacatct	tcattccagct	agctcgggac	ctgcccattg	acacattcac	660

tgtggcaagg	attgacgtgt	ctcagaatga	ccttccttgg	gaatttatgg	tcgatcgtct	720
tcctactgtc	ttgttttttc	cctgcaacag	aaaggaccta	agtgtgaaat	accccgaaga	780
cgccccatc	acccttccaa	acctgttgag	gttcattttg	catcactcag	accctgcttc	840
cagccccag	aatgtggcta	actctcctac	caaggagtgt	cttcagagcg	aggcagtctt	900
acagcggggg	cacatctccc	acttggagag	agagatccag	aaactgagag	cagaaataag	960
cagcctccag	cgagcacaag	tgcagggtgga	gtcccagctc	tccagtgcc	gcagagatga	1020
gcaccggctg	cggcagcagc	agcgggccct	ggaagagcag	cacagcctgc	tccacgcaca	1080
cagtgcagcag	ctgcaggccc	tctatgagca	gaagacacgt	gagctgcagg	agctggccccg	1140
caagctgcag	gagctggccg	atgcctcaga	aaacctcctt	accgagaaca	cgtggctcaa	1200
gatcctggtg	gcgaccatgg	agaggaaact	ggagggcagg	gatggagctg	aaagcctggc	1260
ggcccagaga	gaggtccacc	ccaagcagcc	tgagccctca	gccaccccc	agctccctgg	1320
cagctccctt	ccacctgcca	atgtcagcgc	cacactggtg	tctgaaagga	ataaggagaa	1380
caggacagac	taacttttta	aatgatatga	agaaatcaga	ggtgaaaatt	gtacattggg	1440
aatatattta	tgcaaatttt	attgaaattt	attgtaaata	aagattttct	cagtgggtcta	1500
gaaaatcaaa	aaaaaaaaaa	aaaaaaaaaa	aa			1532

<210> 7

<211> 1532

<212> DNA

<213> Homo sapiens

<400> 7

ttttaacttc	ctatagcccc	ttcagctact	acactgcatg	ttgcaggacc	ataagcaggg	60
gtgtgtcagg	cttcacogac	tctgaacaag	gtgtctttga	agcccctact	gttgcatttt	120
cttcccttga	gaagaaatgt	gaggttgatg	ccccaagctc	cgttcctcac	attgaggaga	180
acaggtatct	ctttccagaa	gtggacatga	ctagcacaaa	cttcacaggc	ctgagctgca	240
gaaccaacaa	gactctcaac	atctaccttt	tggattcaaa	tttgtttggt	tatatgcaga	300
gagactgggt	gctccgagct	ccactcaggt	gaaagaattt	gcggcaattg	ttgacgtgaa	360
agaagaatct	cattacatct	tggatccaaa	gcaagcactg	atgaagctca	ccctagagtc	420
ttttattcaa	aacttcagcg	ttctctatag	tcccttgaaa	aggcatctca	ttggaagtgg	480
ctctgccag	ttcccgctc	agcattttaat	cactgaagtg	acaactgata	ccttttgggg	540
agtagtcctt	caaaaacagg	acgtttctct	gctctattac	gctccgtggt	gcggcttctg	600
tccatccctc	aatcacatct	tcatccagct	agctcggaac	ctgcccattg	acacattcac	660
tgtggcaagg	attgacgtgt	ctcagaatga	ccttccttgg	gaatttatgg	tcgatcgtct	720
tcctactgtc	ttgttttttc	cctgcaacag	aaaggaccta	agtgtgaaat	accccgaaga	780
cgccccatc	acccttccaa	acctgttgag	gttcattttg	catcactcag	accctgcttc	840
cagccccag	aatgtggcta	actctcctac	caaggagtgt	cttcagagcg	aggcagtctt	900
acagcggggg	cacatctccc	acttggagag	agagatccag	aaactgagag	cagaaataag	960
cagcctccag	cgagcacaag	tgcagggtgga	gtcccagctc	tccagtgcc	gcagagatga	1020
gcaccggctg	cggcagcagc	agcgggccct	ggaagagcag	cacagcctgc	tccacgcaca	1080
cagtgcagcag	ctgcaggccc	tctatgagca	gaagacacgt	gagctgcagg	agctggccccg	1140
caagctgcag	gagctggccg	atgcctcaga	aaacctcctt	accgagaaca	cgtggctcaa	1200
gatcctggtg	gcgaccatgg	agaggaaact	ggagggcagg	gatggagctg	aaagcctggc	1260
ggcccagaga	gaggtccacc	ccaagcagcc	tgagccctca	gccaccccc	agctccctgg	1320
cagctccctt	ccacctgcca	atgtcagcgc	cacactggtg	tctgaaagga	ataaggagaa	1380
caggacagac	taacttttta	aatgatatga	agaaatcaga	ggtgaaaatt	gtacattggg	1440
aatatattta	tgcaaatttt	attgaaattt	attgtaaata	aagattttct	cagtgggtcta	1500

gaaaatcaaa aaaaaaaaaa aaaaaaaaaa aa

1532

<210> 8

<211> 759

<212> DNA

<213> Homo sapiens

<400> 8

accactgaga	aaatctttat	ttacaataaa	tttcaataaa	atttgcataa	atatattccc	60
aatgtacaat	tttcacctct	gatttcttca	tatcatttaa	aaagttagtc	tgtcctgttc	120
tccttattcc	tttcagacac	cagtgtggcg	ctgacattgg	caggtggagg	ggagctgcca	180
gggagctggg	gggtggctga	gggctcaggc	tgcttggggg	ggacctctct	ctgggcccgc	240
aggctttcag	ctccatccct	gccctccagt	tccctctcca	tggtcgccac	caggatcttg	300
agccacgtgt	tctcggttaag	gaggttttct	gaggcatcgg	ccagctcctg	cagcttgccg	360
gccagctcct	gcagctcacg	tgtcttctgc	tcatagaggg	cctgcagctg	ctcactgtgt	420
gcgtggagca	ggctgtgctg	ctcttccagg	gcccgcctgt	gctgccgcag	ccggtgctca	480
tctctgcggg	cactggagag	ctgggactcc	acctgcactt	gtgctcgctg	gaggctgctt	540
atttctgctc	tcagtttctg	gatctctctc	tccaagtggg	agatgtgccc	ccgctgtaag	600
actgcctcgc	tctgaagaca	ctccttggtg	ggagagttag	ccacattctg	ggggctggaa	660
agcagggctc	gagtgatgca	aaatgaacct	caacaggttt	ggaaggggtg	tggggacgtc	720
ttcgggggtat	ttcacactta	gggtcctttc	tgttgcagg			759

<210> 9

<211> 775

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (653)..(653)

<223> n = a,t,c, or g

<400> 9

tagaccactg	agaaaatcct	tattttacaat	aaattttcaat	aaaattttgca	taaatatatt	60
cccaatgtac	aatttttcacc	tctgatttct	tcatatcatt	taaaaagtta	gtctgtcctg	120
ttctccttat	tccttttcaga	caccagtgtg	gcgctgacat	tggcaggtgg	aggggagctg	180
ccagggagct	gggggggtggc	tgagggctca	ggctgcttgg	ggtggacctc	tctctggggc	240
gccaggcttt	cagctccatc	cctgccctcc	agtttctctc	ccatggctgc	caccaggatc	300
ttgagccacg	tgttctcggt	aaggaggttt	tctgaggcat	cggccagctc	ctgcagcttg	360
cgggccagct	cctgcagctc	acgtgtcttc	tgctcataga	gggcctgcag	ctgctcactg	420

```

tgtgctgga gcaagctgtg ctgctcttcc agggcccgt gctgctgccg cagccgggtgc 480
tcatctctgc gggcactgga gagctgggac tccacctgca cttgtgctcg ctggaggctg 540
cttatttctg ctctcagttt ctggatctct ctctccaagt gggagatgtg cccccgctgt 600
aagactgcct cgctctgaag acactccttg gtaggagagt tagccacatt ctnggggctg 660
gaagcagggg ctgaatgatg caaaatgaac cctaacaggt ttggaaagggt gatggggagg 720
tcttcggggg tatttcacac ttaggtccct tctgttggac ggaaaaaaaa cagcg 775

```

<210> 10

<211> 533

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (211) .. (211)

<223> n = a,t,g, or c

<400> 10

```

tagaccactg agaaaatctt tatttacaat aaatttcaat aaaatttgca taaatatatt 60
cccaatgtac aattttcacc tctgatttct tcatatcatt taaaaagtta gtctgtcctg 120
ttctccttat tcctttcaga caccagtgtg gcgctgacat tggcaggtgg aggggagctg 180
ccagggagct ggggggtggc tgaggggtca ngctgcttgg ggtggacctc tctctgggcc 240
gccaggcttt cagctccatc cctgccctcc agtttcctct ccatggctgc caccaggatc 300
ttgagccacg tggtctcggt aaggagggtt tctgaggcat cggccagctc ctgcagcttg 360
cgggccagct cctgcagctc acgtgtcttc tgctcataga gggcctgcag ctgctcactg 420
tgtgctgga gcaggtgtg ctgctcttcc agggcccgt gctgctgccg cagccgggtgc 480
tcatctctgc gggcactgga gagctgggac tccacctgca cttgtgctcg ctg 533

```

<210> 11

<211> 543

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (371) .. (371)

<223> n = a,t,g, or c

<220>

<221> misc_feature

<222> (407)..(407)

<223> n = a,t,g, or c

<400> 11

cgagctccac	tcaggtgaaa	gaatttgccg	caattgttga	cgtgaaagaa	gaatctcatt	60
acatcttggg	tccaaagcaa	gcactgatga	agctcaccct	agagtctttt	attcaaaaact	120
tcagcggttct	ctatagtccc	ttgaaaaggc	atctcattgg	aagtggctct	gccagttcc	180
cgtctcagca	tttaatcact	gaagtgacaa	ctgatacctt	ttgggaagta	gtccttcaaa	240
aacaggacgt	tctcctgctc	tattacgctc	cgtgggtgcc	cttctgtcca	tccctcaatc	300
acatcttcat	ccagctagct	cggaaacctgc	ccatggacac	attcactgtg	gcaaggattg	360
acgtgtctca	naatgacctt	ccttggggaat	ttatgggtcga	tctgtctnct	actgtcttgt	420
tttttccctg	caacagaaag	gacctaaagt	tgaaataccc	cgaagacgtc	ctcatcaccc	480
ttccaaacct	ggtgagggtc	attttgcac	actcagaccc	tgttccagc	ccccagatgt	540
ggc						543

<210> 12

<211> 491

<212> DNA

<213> Homo sapiens

<400> 12

tagaccactg	agaaaatctt	tatttacaat	aaatttcaat	aaaatttgca	taaatatatt	60
cccaatgtac	aattttcacc	tctgatttct	tcatatcatt	taaaaagtta	gtctgtcctg	120
ttctccttat	tcctttcaga	caccagtgtg	gcgctgacat	tggcaggtgg	aggggagctg	180
ccagggagct	gggggggtgg	tgagggctca	ggctgcttgg	ggtggacctc	tctctgggcc	240
gccaggcttt	cagctccatc	cctgccctcc	agtttctctc	ccatggctgc	caccaggatc	300
ttgagccacg	tgttctcggt	aaggagggtt	tctgaggcat	cggccagctc	ctgcagcttg	360
cgggccagct	cctgcagctc	acgtgtcttc	tgctcataga	gggcctgcag	ctgctcactg	420
tgtgcgtgga	gcaggtgtgt	ctgctcttcc	agggcccgct	gctgctgccg	cagccgggtgc	480
tcctctctgc	g					491

<210> 13

<211> 488

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (404)..(404)

<223> n=a,t,g, or c

<220>

<221> misc_feature

<222> (406)..(406)

<223> n=a,t,g, or c

<220>

<221> misc_feature

<222> (455)..(455)

<223> n=a,t,g, or c

<400> 13

ctgaagtgac aactgatacc ttttgggaag tagtccttca aaaacaggac gttctcctgc	60
tctattacgc tccgtgggtgc ggcttctgtc catccctcaa tcacatcttc atccagctag	120
ctcggaaacct gcccatggac acattcactg tggcaaggat tgacgtgtct cagaatgacc	180
ttccttggga atttatggtc gatcgtcttc ctactgtctt gttttttccc tgcaacagaa	240
aggacctaag tgtgaaatac cccgaagacg tccccatcac ccttccaaac ctggttgaggt	300
tcattttgca tcactcagac cctgcttcca gccccagaa tgtggctaac tctcctacca	360
aggagtgtct tcagagcgag gcagtcttac agcgggggca catntnccac ttggagagag	420
agatccagaa actgagagca gaaataagca gcctncagcg agcacaagtg caggtggagt	480
cccagctc	488

<210> 14

<211> 468

<212> DNA

<213> Homo sapiens

<400> 14

ctagaccact gagaaaatct ttattttacaa taaattttcaa taaaatttgc ataaatatat	60
tcccaatgta caattttcac ctctgatttc ttcatatcat ttaaaaagtt agtctgtcct	120
gttctcctta ttcttttcag acaccagtgt ggcgctgaca ttggcagggtg gaggggagct	180
gccaggggagc tgggggggtgg ctgaggggtc aggcgtgcttg ggggtggacct ctctctgggc	240
cgccaggctt tcagctccat ccctgccctc cagtttcttc tccatgggtcg ccaccaggat	300


```

cttgagccac gtgttctcgg taaggaggtt ttctgaggca tcggccagct cctgcagctt   360
gcgggccagc tcttgcagct cactgtgtct ctgtcatag agggcctgca gctgctcact   420
gtgtgcgtgg agcaggctgt gctgctcttc cagggcccg cgtgctg   468
<210> 15

```

<211> 459

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (7) .. (7)

<223> n = a,t,g, or c

```

<400> 15
ttttttnatg ttttttttag accactgaga aaatctttat ttacaataaa tttcaataaa   60
atttgcataa atatattccc aatgtacaat tttcacctct gatttcttca tatcatttaa   120
aaagttagtc tgtcctgttc tccttattcc tttcagacac cagtgtggcg ctgacattgg   180
caggtggagg ggagctgcca gggagctggg ggggtggctga gggctcaggc tgcttggggg   240
ggacctctct ctggggccgc aggetttcag ctccatccct gccctccagt ttctcttcca   300
tggtcgccac caggatcttg agccacgtgt tctcggttaag gaggttttct gaggcacggg   360
ccagctcctg cagcttgccg gccagctcct gcagctcacg tgtcttctgc tcatagaggg   420
cctgcagctg ctcaactgtg gcgtggagca ggctgtgct   459
<210> 16

```

<211> 472

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (328) .. (329)

<223> n = a,t,g, or c

```

<400> 16
tctagaccac tgagaaaatc tttattttac ataaatttca ataaaatttg cataaatata   60
ttcccaatgt acaattttca cctctgattt cttcatatca tttaaaagt tagtctgtcc   120

```

```

tggttctcctt attcctttca gacaccagtg tggcgctgac attggcaggt ggaggggagc 180
tgccagggag ctgggggggtg gctgagggct caggctgctt ggggtggacc tctctctggg 240
ccgccacggt ttcagctcca tccctgccct ccagtttctt ctccatgggc gccaccagga 300
tcttgagcca cgtgttctcc gtaagagnnt tttctgaggc atcgccagc tctgcagct 360
tgcgagccag ctctgcagc tcacgtgtct tctgctcata gagggcctgc agctgctcac 420
tgtgtgcgtg gaggcaggctg tgctgctctt ccagggcccg ctgctgctgc cg 472
<210> 17

```

<211> 434

<212> DNA

<213> Homo sapiens

```

<400> 17
tctagaccac tgagaaaatc tttattttaca ataaatttca ataaaatttg cataaatata 60
ttccaatgt acaattttca cctctgattt cttcatatca tttaaaaagt tagtctgtcc 120
tggttctcctt attcctttca gacaccagtg tggcgctgac attggcaggt ggaggggagc 180
tgccagggag ctgggggggtg gctgagggct caggctgctt ggggtggacc tctctctggg 240
ccgccaggct ttcagctcca tccctgccct ccagtttctt ctccatgggc gccaccagga 300
tcttgagcca cgtgttctcg gtaaggaggt tttctgaggc atcgccagc tctgcagct 360
tgcgggccag ctctgcagc tcacgtgtct tctgctcata gagggcctgc agctgctcac 420
tgtgtgcgtg gaggc
<210> 18 434

```

<211> 440

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (218)..(218)

<223> n = a,t,g, or c

<220>

<221> misc_feature

<222> (227)..(227)

<223> n = a,t,g, or c

<220>

<221> misc_feature

<222> (237)..(237)

<223> n = a,t,g, or c

<400> 18

gattttctag accactgaga aaatctttat ttacaataaa tttcaataaa atttgcataa	60
atatattccc aatgtacaat tttcacctct gattttcttca tatcatttaa aaagttagtc	120
tgctctgttc tccttattcc tttcagacac cagtgtggcg ctgacattgg cagggtggagg	180
ggagctgcca gggagctggg ggggtggctga gggctcangc tgcttgnggt ggacctntct	240
ctgggcccgc aggcctttcag ctccatccct gccctccagt ttctctcca tggctgccac	300
caggatcttg agccacgtgt tctcggtgaag gaggttttct gaggcacg cagctcctg	360
cagcttgccg gccagctcct gcagctcacg tgtcttctgc tcatagagg cctgcagctg	420
ctcactgtgt gcgtggagca	440

<210> 19

<211> 446

<212> DNA

<213> Homo sapiens

<400> 19

agaccattta gtaaatcttt atttacaata aatttcaatg aaatttgcat aaatatattc	60
ccaatgtaca attttcacct ctgatttctt catatcattt aaaaagttag tctgtcctgt	120
tctccttatt cttttcagac accagtgtgg cgtgacatt ggcaggtgga ggggagctgc	180
caggagctg gggggtggct gagggctcag gctgcttggg gtggacctct ctctggccg	240
ccagctttc agctccatcc ctgccctcca gtttcctctc catggctgcc accaggatct	300
tgagccacgt gttctcggtg aggaggtttt ctgaggcatc ggccagctcc tgcagcttgc	360
gggccagctc ctgcagctca cgtgtcttct gctcatagag ggcctgcagc tgctcactgt	420
gtgcgtggag caggctgtgc tgctct	446

<210> 20

<211> 440

<212> DNA

<213> Homo sapiens

<400> 20

tttttttttc tagaccactg agaaaatctt tatttacaat aaatttcaat aaaatttgca	60
taaatatatt cccaatgtac aattttcacc tctgatttct tcatatcatt taaaaagtta	120
gtctgtcctg ttctccttat tcctttcaga caccagtgtg gcgctgacat tggcaggtgg	180
aggggagctg ccaggagct ggggggtggc tgagggtcga ggctgcttgg ggtggacctc	240

```

tctctggggc gccacgtttc agctccatcc ctgccctcca gtttcctctc catgggtcgcc 300
accaggatct tgagccacgt gttctcggtg aggagggttt ctgaggcatc ggccagctcc 360
tgcagcttgc gggccagctc ctgcagctca cgtgtcttct gctcatagag ggccctgcagc 420
tgctcactgt gtgcgtggag 440

```

<210> 21

<211> 441

<212> DNA

<213> Homo sapiens

```

<400> 21
tttttttttc tagaccactg agaaaatctt tattttacaat aaattttcaat aaaatttgca 60
taaatatatt cccaatgtac aattttcacc tctgatttct tcatatcatt taaaaagtta 120
gtctgtcctg ttctccttat tcctttcaga caccagtgtg gcgctgacat tggcagggtgg 180
aggggagctg ccaggagct ggggggtggc tgagggtca ggctgcttgg ggtggacctc 240
tctctggggc gccacgttta cagctccatc cctgccctcc agtttcctct ccatggctgc 300
caccaggatc ttgagccacg tgttctcggt aaggagggtt tctgaggcat cggccagctc 360
ctgcagcttg cgggccagct cctgcagctc acgtgtcttc tgctcataga gggcctgcag 420
ctgctcactg tgtgcgtgga g 441

```

<210> 22

<211> 437

<212> DNA

<213> Homo sapiens

```

<400> 22
ctagaccact gagaaaatct ttattttacaa taaatttcaa taaaatttgc ataaatatat 60
tccaatgta caattttcac ctctgatttc ttcatatcat ttaaaaagtt agtctgtcct 120
gttctcctta ttcctttcag acaccagtgt ggcgctgaca ttggcagggtg gaggggagct 180
gccagggagc tggggggtgg ctgagggtc aggetgcttg gggaggacct ctctctgggc 240
cgccacgttt cagctccatc cctgccctcc agtttcctct ccatggctgc caccaggatc 300
ttgagccacg tgttctcggt aaggagggtt tctgaggcat cggccagctc ctgcagcttg 360
cgtgccagct cctgcagctc acgtgtcttc tgctcataga gggcctgcag ctgctcactg 420
tgtgcgtgga gcaggct 437

```

<210> 23

<211> 437

<212> DNA

<213> Homo sapiens

<400> 23

```

ttttctagac cactgagaaa atctttatctt acaataaatt tcaataaaat ttgcataaat    60
atattcccaa tgtacaattt tcacctctga tttcttcata tcatttaaaa agttagtctg    120
tcctgtttctc cttattcctt tcagacacca gtgtggcgct gacattggca ggtggagggg    180
agctgccagg gagctggggg gtggctgagg gctcaggctg cttgggggtg acctctctct    240
gggcgccacc gttatcagcc tccatccctg ccctccagtt tcctctccat ggtcgccacc    300
aggatcttga gccacgtgtt ctcggttaagg aggttttctg aggcacgcgc cagctcctgc    360
agcttgcgag ccagctcctg cagctcacgt gtcttctgct catagagggc ctgcagctgc    420
tcactgtgtg cgtggag
<210> 24

```

<211> 441

<212> DNA

<213> Homo sapiens

```

<400> 24
ttttttctag accactgaga aaatctttat ttacaataaa tttcaataaa atttgcataa    60
atatattccc aatgtacaat tttcacctct gatttcttca tatcatttaa aaagttagtc    120
tgtctgttcc tccttattcc tttcagacac cagtgtggcg ctgacattgg cagggtggacg    180
ggagcagcca gggagctggg ggggtggctga gggctcaggc tgcttggggg ggacctctct    240
ctggggccgc acgtttctag ctccatccct gccctccagt ttcctctcca tggtcgccacc    300
caggatcttg agccacgtgt tctcggttaag gaggttttct gaggcacgcg ccagctcctg    360
cagcttgcgg gccagctcct gcagctcacg tgtcttctgc tcatagaggg gctgaagctg    420
ctcactgtgt gcgtggagaa g
<210> 25

```

<211> 414

<212> DNA

<213> Homo sapiens

```

<400> 25
tttttttttt tttttctaga ccactgagaa aatctttatt tacaataaat ttcaataaaa    60
tttgcataaa tatattccca atgtacaatt ttcacctctg atttcttcat atcatttaaa    120
aagttagtct gtctgtttct ccttattcct ttcagacacc agtgtggcgc tgacattggc    180
agggtggagg gagctgccag ggagctgggg ggtggctgag ggctcaggct gcttgggggtg    240
gacctctctc tggggccgca ggctttcagc tccatccctg ccctccagtt tcctctccat    300
ggtcgccacc aggatcttga gccacgtgtt ctcggttaagg aggttttctg aggcacgcgc    360
cagctcctgc agcttgcggg ccagctcctg cagctcacgt gtcttctgct cata          414
<210> 26

```

<211> 459

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (306)..(306)

<223> n = a,t,g, or c

<400> 26
 tttttttttt tagaccactg agaaaatctt tatttacaat aaatttcaat aaaatttgca 60
 taaatatatt cccaatgtac aattttcacc tctgatttct tcatatcatt taaaaagtta 120
 gtctgtcctg ttctccttat tcctttcaga caccagtgtg gcgctgacat tggcaggtgg 180
 aggggagctg ccaggagct ggggggtggc tgagggtca ggctgcttgg ggtggacctc 240
 tctctgggcc gccaggcttt cagctccatc cctgccctcc agtttcctct ccatggctgc 300
 caccangatc ttgagccacg tgttctcggg agggaggttt tctgaggcat cggccagctc 360
 ctgcagcttg cgggccagct cctgcacttc aacgtgtctt ctgctcatta gagggcctgc 420
 agctgctcac tgtgttgctg ggagcaaggc ctgtgcttg 459
 <210> 27

<211> 399

<212> DNA

<213> Homo sapiens

<400> 27
 ctagaccact gagaaaatct ttattttacaa taaatttcaa taaaatttgc ataaatatat 60
 tccaatgta caattttcac ctctgatttc ttcatatcat ttaaaaagtt agtctgtcct 120
 gttctcctta ttcttttcag acaccagtgt ggcgctgaca ttggcaggtg gaggggagct 180
 gccagggagc tgggggggtgg ctgagggtc aggctgcttg ggggtggacct ctctctgggc 240
 cgccaggctt tcagctccat ccctgccctc cagtttcctc tccatggctg ccaccaggat 300
 cttgagccac gtgttctcgg taaggaggtt ttctgaggca tcggccagct cctgcagctt 360
 gcggggccagc tcctgcagct cacgtgtctt ctgctcata 399
 <210> 28

<211> 350

<212> DNA

<213> Homo sapiens

<400> 28
 atgaagctca ccctagagtc ttttattcaa aacttcagcg ttctctatag tcccttgaaa 60
 aggcattctca ttggaagtgg ctctgcccag ttcccgctctc agcatttaat cactgaagtg 120

```

acaactgata ccttttggga agtagtcctt caaaaacagg acgttctcct gctctattac    180
gctccgtggt gcggcttctg tccatccctc aatcacatct tcatccagct agctcggaac    240
ctgcccattg acacattcac tgtggcaagg attgacgtgt ctcagaatga ccttccttgg    300
gaatttatgg tcgatcgtct tcctactgtc ttgttttttc cctgcaacag              350
<210> 29

```

<211> 116

<212> PRT

<213> Homo sapiens

```

<400> 29
Met Lys Leu Thr Leu Glu Ser Phe Ile Gln Asn Phe Ser Val Leu Tyr
1           5           10           15
Ser Pro Leu Lys Arg His Leu Ile Gly Ser Gly Ser Ala Gln Phe Pro
          20           25           30
Ser Gln His Leu Ile Thr Glu Val Thr Thr Asp Thr Phe Trp Glu Val
          35           40           45
Val Leu Gln Lys Gln Asp Val Leu Leu Leu Tyr Tyr Ala Pro Trp Cys
          50           55           60
Gly Phe Cys Pro Ser Leu Asn His Ile Phe Ile Gln Leu Ala Arg Asn
65           70           75           80
Leu Pro Met Asp Thr Phe Thr Val Ala Arg Ile Asp Val Ser Gln Asn
          85           90           95
Asp Leu Pro Trp Glu Phe Met Val Asp Arg Leu Pro Thr Val Leu Phe
          100          105          110
Phe Pro Cys Asn
          115

```

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 March 2002 (14.03.2002)

PCT

(10) International Publication Number
WO 02/020731 A3

(51) International Patent Classification⁷: **C12N 15/61**,
9/90, 15/63, 5/10, C12P 21/00, C12Q 1/68, 1/533, G01N
33/573, A61K 48/00, A61P 1/00, 5/50, 35/00

(21) International Application Number: PCT/EP01/10125

(22) International Filing Date:
3 September 2001 (03.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/231,297 8 September 2000 (08.09.2000) US

(71) Applicant (for all designated States except US): **BAYER AKTIENGESELLSCHAFT** [DE/DE]; 51368 Leverkusen (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **XIAO, Yonghong** [US/US]; 78 Dana Street #1, Cambridge, MA 02138 (US).

(74) Common Representative: **BAYER AKTIENGESELLSCHAFT**; 51368 Leverkusen (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PI, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CI, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

(88) Date of publication of the international search report:
25 July 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: REGULATION OF HUMAN PROTEIN DISULFIDE ISOMERASE-LIKE ENZYME

(57) Abstract: Reagents which regulate human protein disulfide isomerase-like enzyme and reagents which bind to human protein disulfide isomerase-like enzyme gene products can play a role in preventing, ameliorating, or correcting dysfunctions or disorders including, but not limited to, cancer, diabetes, and obesity.



WO 02/020731 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/10125

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/61 C12N9/90 C12N15/63 C12N5/10 C12P21/00
 C12Q1/68 C12Q1/533 G01N33/573 A61K48/00 A61P1/00
 A61P5/50 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

SEQUENCE SEARCH, EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EM_HUM 'Online! EMBL; 15 March 1999 (1999-03-15) ANDERSSON ET AL.: "Homo sapiens clone 25100 mRNA sequence, complete cds." retrieved from EBI, accession no. AF131780 Database accession no. AF131780 XP002198786 the whole document	1,4,18, 19,25
X	-& DATABASE SWALL 'Online! 1 May 1999 (1999-05-01) ANDERSSON ET AL.: "Hypothetical 37.7 kDa protein" retrieved from EBI, accession no. 095887 Database accession no. 095887 XP002198787 the whole document & ANAL BIOCHEM, vol. 236, 1996, pages 107-113, XP000857646 & GENOME RES, -/--	1,4,18, 19,25



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

14 May 2002

Date of mailing of the international search report

31/05/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 H' Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Ceder, O

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/10125

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	vol. 7, no. 4, 1997, pages 353-358, XP000867830 ---	
X	DATABASE EM_EST 'Online! EMBL; 8 June 2000 (2000-06-08) HEGDE ET AL.: "EST" retrieved from EBI, accession no. AW959209 Database accession no. AW959209 XP002198788 the whole document ---	1,18,19
X	DATABASE EM_EST 'Online! EMBL; 13 March 1999 (1999-03-13) OTTENWALD ET AL.: "EST" retrieved from EBI, accession no. HSM006372 Database accession no. AL048830 XP002198789 the whole document ---	1
A	US 6 001 632 A (BRAXTON SCOTT MICHAEL ET AL) 14 December 1999 (1999-12-14) cited in the application the whole document ---	1-71
A	TASANEN ET AL.: "Promoter of the gene for the multifunctional protein disulfide isomerase polypeptide" THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 267, no. 16, 5 June 1992 (1992-06-05), pages 11513-11519, XP002198785 cited in the application ---	
P,X	WO 01 12790 A (INCYTE GENOMICS INC ;YUE HENRY (US); BANDMAN OLGA (US); BAUGHN MAR) 22 February 2001 (2001-02-22) Seq Id Nos 8 and 16 abstract; claims ---	1-71
P,X	DATABASE EM_HUM 'Online! EMBL; 15 May 2001 (2001-05-15) ISOGAI ET AL.: "Homo sapiens cDNA FLJ14558 fis" retrieved from EBI, accession no. AK027464 Database accession no. AK027464 XP002198790 the whole document ---	1,4,18, 19,25
T	-& DATABASE SWALL 'Online! 1 December 2001 (2001-12-01) ISOGAI ET AL.: "cDNA FLJ14558 fis" retrieved from EBI, accession no. Q96K53 Database accession no. Q96K53 XP002198791 the whole document -----	1,4,18, 19,25

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claims 68-71 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Continuation of Box I.2

Present claims 13-17, 60, 62 and 68-71 relate to compounds/methods defined by reference to desirable characteristics or properties, namely binding to any polynucleotide of claim 1, or to any polypeptide of claim 4, or being identified by the method of claim 10-12, or modulating the function of a human protein disulfide isomerase-like enzyme.

The claims cover all compounds/methods having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compounds/methods. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the compound/method by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the compound being antibody to the polypeptides of claim 4 or being polynucleotides of claim 1.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/10125

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6001632	A	14-12-1999	US 5798249 A	25-08-1998
			AU 3066197 A	05-12-1997
			WO 9743426 A1	20-11-1997
<hr/>				
WO 0112790	A	22-02-2001	AU 6911300 A	13-03-2001
			WO 0112790 A2	22-02-2001
<hr/>				

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)